



Heteroradulum yunnanensis sp. nov. (Auriculariales, Basidiomycota) evidenced by morphological characters and phylogenetic analyses in China

QIAN-XIN GUAN^{1,2}, CHAO-MAO LIU², TANG-JIE ZHAO³ & CHANG-LIN ZHAO^{1,2,4*}

¹Key Laboratory for Forest Resources Conservation and Utilization in the Southwest Mountains of China, Ministry of Education, Southwest Forestry University, Kunming 650224, P.R. China

²College of Biodiversity Conservation, Southwest Forestry University, Kunming 650224, P.R. China

³Wenshan Forestry Research Institute, Wenshan, Yunnan 663300, P.R. China

⁴Key Laboratory of Forest Disaster Warning and Control of Yunnan Province, Southwest Forestry University, Kunming 650224, P.R. China

*Corresponding author's e-mail: fungichanglinz@163.com

Abstract

A new wood-inhabiting fungal species, *Heteroradulum yunnanensis*, is proposed based on a combination of morphological features and molecular evidence. The species is characterized by an annual growth habit, resupinate basidiomata with odontoid hymenial surface (50–100 µm long), more or less pronounced yellow stains in older basidiomata, a monomitic hyphal system with thin-walled, clamped generative hyphae and two to three-celled basidia and cylindrical, hyaline, thin-walled, smooth, IKI–, CB– basidiospores measuring as 17–24 × 5–8 µm. Sequences of ITS and LSU nrRNA gene regions of the studied samples were generated, and phylogenetic analyses were performed with maximum likelihood, maximum parsimony and bayesian inference methods. The phylogenetic analyses based on molecular data of ITS+nLSU sequences showed that *Heteroradulum yunnanensis* formed a monophyletic lineage with a strong support (100% BS, 100% BP, 1.00 BPP) and then grouped with *H. adnatum*.

Keywords: *Exidiopsis*, phylogeny, taxonomy, wood-inhabiting fungi, Yunnan Province

Introduction

Auriculariales Bromhead is a special wood-inhabiting fungal order with high morphological diversity, and it not only includes jelly species (Wu *et al.* 2015a, b), but many species that are previously called corticioid fungi belong to this order. *Heteroradulum* Lloyd ex Spirin & Malysheva (2017: 709) is one of the genera with corticioid hymenophore. The genus was typified by *H. kmetii* (Bres.) Spirin & Malysheva (2017: 711) (Malysheva & Spirin 2017), which is characterized by a combination of annual or perennial, resupinate or effused-reflexed basidiomata with a leathery consistence, hymenophore smooth to odontoid, a monomitic hyphal structure (rare dimitic) with clamp connections on generative hyphae, and the presence of cystidia, basidia narrowly ovoid to obconical, longitudinally septate with a well-developed enucleate stalk and basidiospores hyaline, thin-walled, smooth, cylindrical, acyanophilous and negative in Melzer's reagent (Malysheva & Spirin 2017). So far six species have been accepted in the genus worldwide (Berkeley 1873, Berkeley & Broome 1875, Patouillard & Lagerheim 1893, Bresadola 1897, Bodman 1953, Malysheva & Spirin 2017).

Recently, molecular studies involving *Heteroradulum* have been carried out (Malysheva & Spirin 2017, Malysheva *et al.* 2018, Yuan *et al.* 2018). Phylogenetic study of the Auriculariales with stereoid basidiocarps revealed that *Heteroradulum* was validated and two new species, five combinations were introduced on the base of the combined data of the nuclear ribosomal LSU and the internal transcribed spacer (ITS) regions (Malysheva & Spirin 2017). Malysheva *et al.* (2018) revised of *Protohydnum* A. Möller (1895:173) and suggested that *Heteroradulum* grouped with *Exidiopsis* (Bref.) Möller (1895:85) and *Tremellochaete* Raitv (1964:29). Yuan *et al.* (2018) employed molecular study based on ITS+nLSU datasets to study of *Grammatus* H.S. Yuan & Decock (2018:32) in Auriculariales from tropical China and revealed that *Heteroradulum* grouped with *Exidiopsis* and *Grammatus*.

During investigations on wood-inhabiting fungi in southern China, an additional taxon was found which could

not be assigned to any described species. In this study, the authors expand samplings from previous studies to examine taxonomy and phylogeny of this new species within the *Heteroradulum*, based on the internal transcribed spacer (ITS) regions and the large subunit nuclear ribosomal RNA gene (nLSU) sequences.

TABLE 1. List of species, specimens, and GenBank accession numbers of sequences used in this study.

Species name	Sample no.	GenBank accession no.		References
		ITS	nLSU	
<i>Exidiopsis effusa</i>	OM 19136	KX262145	KX262193	Malysheva & Spirin 2017
<i>Heteroradulum adnatum</i>	LR 23453	KX262116	KX262165	Malysheva & Spirin 2017
<i>H. deglubens</i>	LE 38182	KX262112	KX262162	Malysheva & Spirin 2017
<i>H. deglubens</i>	Solheim 1864	KX262133	KX262181	Malysheva & Spirin 2017
<i>H. kmetii</i>	Kmet	KX262124	KX262173	Malysheva & Spirin 2017
<i>H. kmetii</i>	OF 295640	KX262122	KX262171	Malysheva & Spirin 2017
<i>H. kmetii</i>	LR 33201	KX262117	KX262166	Malysheva & Spirin 2017
<i>H. kmetii</i>	OF 295639	KX262128	KX262177	Malysheva & Spirin 2017
<i>H. yunnanensis</i>	CLZhao 4023	MT215564	MT215568	This study
<i>H. yunnanensis</i>	CLZhao 8106	MT215565	MT215569	This study
<i>H. yunnanensis</i>	CLZhao 9132	MT215566	MT215570	This study
<i>H. yunnanensis</i>	CLZhao 9200	MT215567	MT215571	This study
<i>Tremellochaete japonica</i>	LE 303446	KX262110	KX262160	Malysheva & Spirin 2017

Materials and methods

Morphological studies.—The specimens studied are deposited at the herbarium of Southwest Forestry University (SWFC), Kunming, Yunnan Province, P.R. China. Macro-morphological descriptions are based on field notes. Special colour terms follow Petersen (1996). Micro-morphological data were obtained from the dried specimens, and observed under a light microscope following Dai (2012). The following abbreviations were used: KOH = 5% potassium hydroxide, CB = Cotton Blue, CB– = acyanophilous, IKI = Melzer’s reagent, IKI– = both inamyloid and indextrinoid, L = mean spore length (arithmetic average of all spores), W = mean spore width (arithmetic average of all spores), Q = variation in the L/W ratios between the specimens studied, n (a/b) = number of spores (a) measured from given number (b) of specimens.

Molecular procedures and phylogenetic analyses.—CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd, Beijing) was used to obtain genomic DNA from dried specimens, according to the manufacturer’s instructions with some modifications that a small piece of dried fungal specimen (about 30 mg) was ground to powder with liquid nitrogen. The powder was transferred to a 1.5 mL centrifuge tube, suspended in 0.4 mL of lysis buffer, and incubated in a 65 °C water bath for 60 min. After that, 0.4 mL phenol-chloroform (24:1) was added to each tube and the suspension was shaken vigorously. After centrifugation at 13,000 rpm for 5 min, 0.3 mL supernatant was transferred to a new tube and mixed with 0.45 mL binding buffer. The mixture was then transferred to an adsorbing column (AC) for centrifugation at 13,000 rpm for 0.5 min. Then, 0.5 mL inhibitor removal fluid was added in AC for a centrifugation at 12,000 rpm for 0.5 min. After washing twice with 0.5 mL washing buffer, the AC was transferred to a clean centrifuge tube, and 100 ml elution buffer was added to the middle of adsorbed film to elute the genome DNA. ITS region was amplified with primer pairs ITS5 and ITS4 (White *et al.* 1990). Nuclear LSU region was amplified with primer pairs LR0R and LR7 (<http://www.biology.duke.edu/fungi/mycolab/primers.htm>). The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min, followed by 35 cycles at 94 °C for 40 s, 58 °C for 45 s and 72 °C for 1 min, and a final extension of 72 °C for 10 min. The PCR procedure for nLSU was as follows: initial denaturation

at 94 °C for 1 min, followed by 35 cycles at 94 °C for 30 s, 48 °C 1 min and 72 °C for 1.5 min, and a final extension of 72 °C for 10 min. The PCR products were purified and directly sequenced at Kunming Tsingke Biological Technology Limited Company. All newly generated sequences were deposited at GenBank (Table 1).

Sequencher 4.6 (GeneCodes, Ann Arbor, MI, USA) was used to edit the DNA sequence. Sequences were aligned in MAFFT 6 (Kato & Toh 2008, <http://mafft.cbrc.jp/alignment/server/>) using the “G-INS-I” strategy and manually adjusted in BioEdit (Hall 1999). The sequence alignment was deposited in TreeBase (submission ID 25001). Sequences of *Exidiopsis effusa* (Bref. ex Sacc.) Möller (1895:168) and *Tremellochaete japonica* (Lloyd) Raitv (1964:30), obtained from GenBank were used as outgroups to root trees following Malysheva & Spirin (2017) in the ITS+nLSU analysis (Fig. 1).




FIGURE 1. Maximum parsimony strict consensus tree illustrating the phylogeny of *Heteroradulum yunnanensis* and related species in *Heteroradulum* based on ITS+nLSU sequences. Branches are labeled with maximum likelihood bootstrap higher than 70%, parsimony bootstrap proportions higher than 50% and Bayesian posterior probabilities more than 0.95 respectively.

Maximum parsimony analysis was applied to the ITS+nLSU dataset sequences. Approaches to phylogenetic analysis followed Chen *et al.* (2016), and the tree construction procedure was performed in PAUP* version 4.0b10 (Swofford 2002). All characters were equally weighted and gaps were treated as missing data. Trees were inferred using the heuristic search option with TBR branch swapping and 1000 random sequence additions. Max-trees were set to 5000, branches of zero length were collapsed and all parsimonious trees were saved. Clade robustness was assessed using a bootstrap (BT) analysis with 1,000 replicates (Felsenstein 1985). Descriptive tree statistics tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC), and homoplasy index (HI) were calculated for each Maximum Parsimonious Tree generated. Sequences were also analyzed using Maximum Likelihood (ML) with RAxML-HPC2 through the Cipres Science Gateway (www.phylo.org, Miller *et al.* 2009). Branch support (BS) for ML analysis was determined by 1000 bootstrap replicate.




FIGURE 2. Basidiomata of *Heteroradulum yunnanensis* (holotype). Bars: A = 2 cm, B = 2 mm.

MrModeltest 2.3 (Nylander 2004) was used to determine the best-fit evolution model for each data set for Bayesian inference (BI). Bayesian inference was calculated with MrBayes 3.1.2 with a general time reversible (GTR+G) model of DNA substitution and a gamma distribution rate variation across sites (Ronquist & Huelsenbeck 2003). Four Markov chains run for 2 runs from random starting trees for 40, 000 generations (Fig. 1) and trees were sampled every 100 generations. The first one-fourth generations were discarded as burn-in. A majority rule consensus tree of all remaining trees was calculated. Branches were considered as significantly supported if they received maximum likelihood bootstrap (ML) >70%, maximum parsimony bootstrap (MP) >50%, or Bayesian posterior probabilities (PP) >0.95.




FIGURE 3. Microscopic structures of *Heteroradulum yunnanensis* (drawn from the holotype). a. Basidiospores. b. Basidia. c. Cystidia. d. A section of hymenium. Bars: a,b,c,d = 10 μ m.

TABLE 2. A morphological comparison between *Heteroradulum yunnanensis* and four similar species in the genus.

Taxon	Habit	Hymenial surface	Hyphal system	Basidia (μm)	Basidiospores (μm)	Cystidia (μm)
<i>Heteroradulum adnatum</i>	Annual	White	Monomitic	17.3–21.8 × 9.2–11.3	11.4–14.2 × 5.2–7.2	26–95 × 4–10
<i>H. deglubens</i>	Annual or perennial	Greyish pink	Monomitic	26–44.8 × 9.3–14	13–20.4 × 6–8.2	31–44 × 5–6
<i>H. kmetii</i>	Perennial	Pinkish grey	Dimitic	27.4–45.4 × 9.7–14.7	14.1–24.9 × 5.8–9.7	27–40 × 4–7.5
<i>H. spinulosum</i>	Annual	Ochraceous to pinkish grey	Dimitic	17.4–29.8 × 9.2–12.4	11.7–19.8 × 5.9–7.4	23–45 × 6–7
<i>H. yunnanensis</i>	Annual	White	Monomitic	28–41 × 9–14	15–24.5 × 4.5–8.5	13–35 × 2–6

Results

Molecular phylogeny

The ITS+nLSU dataset (Fig. 1) included sequences from 13 fungal specimens representing six species. The dataset had an aligned length of 1971 characters, of which 1793 characters are constant, 86 are variable and parsimony-uninformative, and 92 are parsimony-informative. Maximum parsimony analysis yielded 6 equally parsimonious trees (TL = 236, CI = 0.898, HI = 0.101, RI = 0.916, RC = 0.823). Best model for the ITS+nLSU dataset estimated and applied in the Bayesian analysis: GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet (1,1,1,1). Bayesian analysis and ML analysis resulted in a similar topology as MP analysis, with an average standard deviation of split frequencies = 0.009361 (BI).

The phylogenetic tree (Fig. 1) inferred from ITS+nLSU sequences, demonstrated four species of *Heteroradulum* and revealed that the new species formed a monophyletic lineage with high supports of 100% BS, 100% BP and 1.00 BPP and then grouped with *H. adnatum* Spirin & Malysheva.

Taxonomy

Heteroradulum yunnanensis C.L. Zhao, *sp. nov.* (Figs. 2, 3)

Mycobank no.: MB 834997

Holotype.—China. Yunnan Province: Yuxi, Xinping County, Jinshan Forestry Park, on a fallen angiosperm branch, 2 January 2019, *CLZhao 9132* (SWFC).

Etymology.—*Yunnanensis* (Lat.): referring to the locality (Yunnan) of the type specimens.

Description.—Basidiomata annual, resupinate, leathery, without odour or taste when fresh, becoming membranaceous upon drying, up to 15 cm long, 100–300 µm thick. Hymenial surface odontoid, 50–100 µm long, white to slightly grey when fresh, becoming white to smoke grey upon drying, older basidiomata with more or less pronounced yellow stains. Sterile margin smoke grey. Hyphal system monomitic; generative hyphae with clamp connections, IKI–, CB–; tissues unchanged in KOH; subiculum almost absent or indistinct; hymenium thickening, hyphae colorless, more or less interwoven, thin-walled, unbranched, 2–4 µm in diameter.

Hymenium.—Cystidia clavate to fusiform, 13–35 × 2–6 µm; basidia narrowly ovoid to obconical, longitudinally septate, two to three-celled, embedded, often with a well-developed enucleate stalk, 28–41 × 9–14 µm. Basidiospores hyaline, thin-walled, smooth, cylindrical sometimes curved, with oily inclusions, IKI–, CB–, (15–)17–24(–24.5) × (4.5–)5–8(–8.5) µm, L = 20.6 µm, W = 6.71 µm, Q = 2.63–3.42 (n = 90/3).

Ecology and distribution.—Lignicolous, causing a white rot. Found in China.

Additional specimens (paratypes) examined.—China. Yunnan Province: Yuxi, Xinping County, Jishan Forestry Park, on a fallen angiosperm branch, 2 January 2019, *CLZhao 9200* (SWFC), 21 August 2018, *CLZhao 8106* (SWFC); Puer, Jingdong County, Ailaoshan National Nature Reserve, on the fallen angiosperm branch, 4 October 2017, *CLZhao 4023* (SWFC).

Discussion

In the present study, a new species, *Heteroradulum yunnanensis*, is described based on phylogenetic analyses and morphological characters.

Previously, seven corticioid genera in the Auriculariales were phylogenetically studied: *Amphistereum* Spirin & Malysheva (2017:696), *Eichleriella* Bres. (1903:115), *Exidiopsis*, *Heteroradulum*, *Hirneolina* (Pat.) Bres. (1905:208), *Sclerotrema* Spirin & Malysheva (2017:712) and *Tremellochaete* (Malysheva & Spirin 2017). Our result based on the combined ITS+nLSU sequence data (Fig. 1) demonstrated that *H. yunnanensis* was nested in the *Heteroradulum*.

Phylogenetically, *Heteroradulum yunnanensis* is closely related to *H. adnatum* in the rDNA based on the phylogeny (Fig. 1). But morphologically *H. adnatum* differs from *H. yunnanensis* by having the pink hymenial surface and smaller basidiospores (11.4–14.2 × 5.2–7.2 µm, Malysheva & Spirin 2017).

Morphologically, *Heteroradulum yunnanensis* resembles four similar species in this genus, *H. adnatum*, *H. deglubens* (Berk. & Broome) Spirin & Malysheva (2017:710), *H. kmetii* and *H. spinulosum* (Berk. & M.A. Curtis) Spirin & Malysheva (2017:712). A morphological comparison between *H. yunnanensis* and four species is presented in Table 2.

Wood-rotting fungi are an extensively studied group of Basidiomycota (Gilbertson & Ryvarden 1986, 1987, Núñez & Ryvarden 2001, Bernicchia & Gorjón 2010, Dai 2012, Ryvarden & Melo 2014, Wu *et al.* 2017, Shen *et al.* 2019), because some of them are economically important (Dai *et al.* 2007, Wu *et al.* 2019a), but the Chinese wood-rotting fungi diversity is still not well known, especially in subtropics and tropics, many recently described taxa of wood-rotting fungi were from these areas (Zhao & Wu 2017, Luo *et al.* 2018, Shen *et al.* 2018, Liu *et al.* 2019, Wu *et al.* 2019b). The new species in the present study, *Heteroradulum yunnanensis*, is from subtropics, too. It is possible that new taxa will be found after further investigations and molecular analyses, especially in some poorly known groups.

Acknowledgements

The research is supported by the National Natural Science Foundation of China (Project No. 31700023) and Yunnan Agricultural Foundation Projects (2017FG001-042) and the Science Foundation of Southwest Forestry University (Projects No. 111715, QN201904).

References

- Berkeley, M.J. (1873) Notices of North American fungi. *Grevillea* 1: 145–150.
<https://doi.org/10.2307/1218613>
- Berkeley, M.J. & Broome, C.E. (1875) Notices of British fungi (1402-1500). *Annual Magazine Natural History Series* 15: 28–41.
<https://doi.org/10.1080/00222937508681018>
- Bernicchia, A. & Gorjón, S.P. (2010) *Fungi Europaei 12: Corticiaceae s.l.* Edizioni Candusso, Lomazzo, pp. 1–1007.
- Bodman, M.C. (1953) A taxonomic study of the genus *Heterochaete*. *Lloydia* 15: 193–233.
- Bresadola, G. (1897) Hymenomycetes Hungarici Kmetiani. *Atti dell'Istituto Reale dell'Accademia di Rovereto di Scienze* 3: 66–120.
- Bresadola, G. (1903) Fungi Polonici a cl. Viro B. Eichler lecti. *Annales Mycologici* 1: 97–131.
- Chen, J.J., Cui, B.K. & Dai, Y.C. (2016) Global diversity and molecular systematics of *Wrightoporia* s. l. (Russulales, Basidiomycota). *Persoonia* 37: 21–36.
<https://doi.org/10.3767/003158516X689666>
- Dai, Y.C. (2012) Polypore diversity in China with an annotated checklist of Chinese polypores. *Mycoscience* 53: 49–80.
<https://doi.org/10.1007/s10267-011-0134-3>
- Dai, Y.C., Cui, B.K., Yuan, H.S. & Li, B.D. (2007) Pathogenic wood-decaying fungi in China. *Forest Pathology* 37: 105–120.
<https://doi.org/10.1111/j.1439-0329.2007.00485.x>
- Felsenstein, J. (1985) Confidence intervals on phylogenetics: an approach using bootstrap. *Evolution* 39: 783–791.
<https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
- Gilbertson, R.L. & Ryvarden, L. (1986–1987) North American polypores 1-2. *Fungiflora*, Oslo. pp. 1–885.
- Hall, T.A. (1999) Bioedit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41: 95–98.
- Katoh, K. & Toh, H. (2008) Recent developments in the MAFFT multiple sequence alignment program. *Briefings Bioinform* 9: 286–298.
<https://doi.org/10.1093/bib/bbn013>
- Liu, X.F., Shen, S. & Zhao, C.L. (2019) Morphological and molecular identification of a new species of *Eichleriella* (Auriculariales, Basidiomycota) in China. *Phytotaxa* 404: 245–254.
<https://doi.org/10.11646/phytotaxa.404.6.3>
- Luo, K.Y., Ma, X. & Zhao, C.L. (2019) *Neofavolus yunnanensis* sp. nov. (Polyporales, Basidiomycota) from China: evidence from morphology and DNA sequence data. *Phytotaxa* 408: 109–116.
<https://doi.org/10.11646/phytotaxa.408.2.3>
- Malysheva, V. & Spirin, V. (2017) Taxonomy and phylogeny of the Auriculariales (Agaricomycetes, Basidiomycota) with stereoid basidiocarps. *Fungal Biology* 121: 689–715.
<https://doi.org/10.1016/j.funbio.2017.05.001>
- Malysheva, V., Spirin, V., Miettinen, O., Motato-Vásquez, Viviana, Hernawati & Seelan, J.S.S. (2018) Revision of *Protohydnum*

- (Auriculariales, Basidiomycota). *Mycological Progress* 17: 805–814.
<https://doi.org/10.1007/s11557-018-1393-6>
- Miller, M.A., Holder, M.T., Vos, R., Midford, P.E., Liebowitz, T., Chan, L., Hoover, P. & Warnow, T. (2009) The CIPRES Portals. CIPRES. Available from: http://www.phylo.org/sub_sections/portal (accessed 4 August 2009) [Archived by WebCite(r) at: <http://www.webcitation.org/5imQlJeQa>]
- Möller, A. (1895) Protobasidiomyceten. *Botanische Mitteilungen aus den Tropen* 8: 1–180.
<https://doi.org/10.1093/bioinformatics/btg180>
- Núñez, M. & Ryvarden, L. (2001) East Asian polypores 2. *Synopsis Fungorum* 14: 165–522.
- Nylander, J.A.A. (2004) *MrModeltest v2*. Program distributed by the author. Evolutionary Biology Centre, Uppsala University.
- Patouillard, N.T. & Lagerheim, G.de. (1893) Champignons de l'Equateur (Pugillus III). *Bulletin de la Société Mycologique de France* 9: 124–165.
- Petersen, J.H. (1996) *Farvekort. The Danish Mycological Society's colour-chart*. Foreningen til Svampekundskabens Fremme, Greve, pp. 1–6.
- Raitviir, A. (1964) Some new species and a new genus of *Discomycetes* and Heterobasidiomycetidae. *Eesti NSV Teaduste Akadeemia Toimetised* 13: 28–32.
- Ronquist, F. & Huelsenbeck, J.P. (2003) MrBayes 3: bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574.
<https://doi.org/10.1093/bioinformatics/btg180>
- Ryvarden, L. & Melo, I. (2014) Poroid fungi of Europe. *Synopsis Fungorum* 31: 1–455.
- Saccardo, P.A. & Saccardo, D. (1905) Supplementum universale. Pars VI. Hymenomycetae-Laboulbeniomycetae. *Sylloge Fungorum* 17: 1–991.
- Shen, L.L., Wang, M., Zhou, J.L., Xing, J.H., Cui, B.K. & Dai, Y.C. (2019) Taxonomy and phylogeny of *Postia*. Multi-gene phylogeny and taxonomy of the brown-rot fungi: *Postia* (Polyporales, Basidiomycota) and related genera. *Persoonia* 42: 101–126.
<https://doi.org/10.3767/persoonia.2019.42.05>
- Shen, S., Ma, X., Xu, T.M. & Zhao, C.L. (2018) *Phlebia ailaoshanensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analyses. *Phytotaxa* 373: 184–196.
<https://doi.org/10.11646/phytotaxa.373.3.2>
- Swofford, D.L. (2002) *PAUP*: phylogenetic analysis using parsimony (*and other methods)*. Version 4.0b10. Sinauer Associates, Massachusetts.
- White, T.J., Bruns, T., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis, M.A., Gelfand, D.H., Sninsky, J.J. & White, T.J. (Eds.) *PCR protocols: a guide to methods and applications*. Academic Press, San Diego, pp. 315–322.
<https://doi.org/10.1016/B978-0-12-372180-8.50042-1>
- Wu, F., Yuan, Y., He, S.H., Bandara, A.R., Hyde, K.D., Malysheva, V.F. & Dai, Y.C. (2015a) Global diversity and taxonomy of the *Auricularia auricula-judae* complex (Auriculariales, Basidiomycota) *Mycological Progress* 14: 95.
<https://doi.org/10.1007/s11557-015-1113-4>
- Wu, F., Yuan, Y., Rivoire, B. & Dai, Y.C. (2015b) Phylogeny and diversity of the *Auricularia mesenterica* (Auriculariales, Basidiomycota) complex. *Mycological Progress* 14: 42.
<https://doi.org/10.1007/s11557-015-1065-8>
- Wu, F., Chen, J.J., Ji, X.H., Vlasák, J. & Dai, Y.C. (2017) Phylogeny and diversity of the morphologically similar polypore genera *Rigidoporus*, *Physisporinus*, *Oxyporus*, and *Leucophellinus*. *Mycologia* 109: 749–765.
<https://doi.org/10.1080/00275514.2017.1405215>
- Wu, F., Zhou, L.W., Yang, Z.L., Bau, T., Li, T.H. & Dai, Y.C. (2019a) Resource diversity of Chinese macrofungi: edible, medicinal and poisonous species. *Fungal Diversity* 98: 1–76.
<https://doi.org/10.1007/s13225-019-00432-7>
- Wu, Y.X., Shen, S. & Zhao, C.L. (2019b) *Podoscypha yunnanensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analyses. *Phytotaxa* 287: 210–218.
<https://doi.org/10.11646/phytotaxa.387.3.2>
- Yuan, H.S., Lu, X. & Decock, C. (2018) Molecular and morphological evidence reveal a new genus and species in Auriculariales from tropical China. *MycKeys* 35: 27–39.
<https://doi.org/10.3897/mycokeys.35.25271>
- Zhao, C.L. & Wu, Z.Q. (2017) *Ceriporiopsis kunmingensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. *Mycological Progress* 16: 93–100.
<https://doi.org/10.1016/j.myc.2013.11.006>