





A peer-reviewed open-access journal MycooKeys Launched to accelerate biodiversity research

# Morphological and phylogenetic evidence for recognition of two new species of Hyphoderma (Basidiomycota) from southern China, with a key to all Chinese Hyphoderma

**RESEARCH ARTICLE** 

Qian-Xin Guan<sup>1,3</sup>, Yi-Fei Li<sup>3</sup>, Chang-Lin Zhao<sup>1,2,3</sup>

I Key Laboratory for Forest Resources Conservation and Utilization in the Southwest Mountains of China, Ministry of Education, Southwest Forestry University, Kunming 650224, China 2 Yunnan Academy of Biodiversity, Southwest Forestry University, Kunming 650224, China 3 College of Biodiversity Conservation, Southwest Forestry University, Kunming 650224, China

Corresponding author: Chang-Lin Zhao (fungi@swfu.edu.cn)

Academic editor: R. Henrik Nilsson | Received 9 June 2021 | Accepted 28 August 2021 | Published 20 September 2021

**Citation:** Guan Q-X, Li Y-F, Zhao C-L (2021) Morphological and phylogenetic evidence for recognition of two new species of *Hyphoderma* (Basidiomycota) from southern China, with a key to all Chinese *Hyphoderma*. MycoKeys 83: 145–160. https://doi.org/10.3897/mycokeys.83.69909

### Abstract

Wood-inhabiting fungi play crucial roles as decomposers in forest ecosystems and, in this study, two new wood-inhabiting corticioid fungi, *Hyphoderma puerense* and *H. tenuissimum* **spp. nov.**, are proposed, based on a combination of morphological features and molecular evidence. *Hyphoderma puerense* is characterised by effused basidiomata with smooth to floccose hymenial surface, a monomitic hyphal system with clamped generative hyphae and ellipsoid basidiospores. *Hyphoderma tenuissimum* is characterised by resupinate basidiomata with tuberculate to minutely-grandinioid hymenial surface, septate cystidia and cylindrical to allantoid basidiospores. Sequences of ITS and nLSU rRNA markers of the studied samples were generated and phylogenetic analyses were performed with Maximum Likelihood, maximum parsimony and Bayesian Inference methods. These analyses showed that the two new species clustered into *Hyphoderma*, in which *H. puerense* grouped with *H. moniliforme* and *H. tenuissimum* formed a singleton lineage. In addition, an identification key to Chinese *Hyphoderma* is provided.

### Keywords

Corticioid fungi, diversity, Hyphodermataceae, molecular phylogeny, taxonomy, Yunnan Province

### Introduction

Fungi are eukaryotic microorganisms that play fundamental ecological roles as decomposers and mutualists of plants and animals. They drive carbon cycling in forest soils, mediate mineral nutrition of plants and alleviate carbon limitations of other soil organisms (Tedersoo et al. 2014). Fungi form an ecologically important branch of the tree of life, based on their distinct and diverse characters (James et al. 2020).

*Hyphoderma* Wallr. was typified by *H. setigerum* (Fr.) Donk (Donk 1957) and the genus is characterised by resupinate to effuse-reflexed basidiomata of ceraceous consistency and a smooth to tuberculate or hydnoid hymenophore. *Hyphoderma* species are characterised by a monomitic (rarely dimitic) hyphal structure with clamp connections on generative hyphae, presence of cystidia or not, suburniform to subcylindrical to cylindrical basidia and ellipsoid to subglobose, smooth, thin-walled basidiospores (Wallroth 1833; Bernicchia and Gorjón 2010). Currently, about 105 species have been accepted in *Hyphoderma* worldwide (Donk 1957; Nakasone 2008; Wu et al. 2010; Baltazar et al. 2016; Martín et al. 2018; Guan and Zhao 2021a, 2021b; Ma et al. 2021). Index Fungorum (http://www.indexfungorum.org; accessed on 16 July 2021) and MycoBank (https://www.mycobank.org; accessed on 16 July 2021) register 199 specific and infraspecific names in *Hyphoderma*.

*Hyphoderma* has been studied using molecular data, particularly the internal transcribed spacer (ITS) region and the large subunit nuclear ribosomal RNA gene (nLSU). Larsson (2007) showed that *H. obtusum* J. Erikss. and *H. setigerum* clustered into the Meruliaceae Rea and formed a sister taxon to *Hypochnicium polonense* (Bres.) Å. Strid. Telleria et al. (2012) proposed a new species, *Hyphoderma macaronesicum* Tellería, M. Dueñas, Beltrán-Tej., Rodr.-Armas & M.P. Martín and then discussed the relationships with the closely-related taxa in *Hyphoderma*. Research into the *Hyphoderma setigerum* complex showed that *H. pinicola* Yurch. & Sheng H. Wu represented a fifth species in this complex (Yurchenko and Wu 2014b). A revised family-level classification of the Polyporales revealed that four *Hyphoderma* species grouped into the residual polyporoid clade, belonging to Hyphodermataceae in that they grouped with three related genera in Meripilaceae: *Meripilus* P. Karst., *Physisporinus* P. Karst. and *Rigidoporus* Murrill (Justo et al. 2017).

In this study, two undescribed species of corticioid fungi from forest ecosystems were collected in the Yunnan Province, China. We present morphological and molecular phylogenetic evidence that support the recognition of two new species in *Hyphoderma*, based on the nuclear ribosomal internal transcribed spacer region (ITS1, 5.8S and ITS2) and the nuclear ribosomal nLSU (28S) gene.

### Materials and methods

#### Morphology

The studied specimens are deposited at the Herbarium of Southwest Forestry University (SWFC), Kunming, Yunnan Province, P.R. China. Macromorphological descriptions are based on field notes and photos captured in the field and lab. Colour terminology fol-

lows Petersen (Petersen 1996). Micromorphological data were obtained from the dried specimens when observed under a light microscope following Dai (2012). The following abbreviations are used: **KOH** = 5% potassium hydroxide water solution, **CB** = Cotton Blue, **CB**– a acyanophilous, **IKI** = Melzer's Reagent, **IKI**– = both inamyloid and indextrinoid, **L** = mean spore length (arithmetic average for all spores), **W** = mean spore width (arithmetic average for all spores), **Q** = variation in the L/W ratios between the specimens studied and **n** = a/b (number of spores (a) measured from given number (b) of specimens).

## Molecular phylogeny

The CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd, Beijing) was used to obtain genomic DNA from the dried specimens following the manufacturer's instructions (as done in Zhao and Wu 2017). The nuclear ribosomal ITS region was amplified with the primers ITS5 and ITS4 (White et al. 1990). The nuclear ribosomal LSU gene was amplified with the primers LR0R and LR7 (Vilgalys and Hester 1990; Rehner and Samuels 1994). The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min followed by 35 cycles at 94 °C for 40 s, 58 °C for 45 s and 72 °C for 1 min and a final extension of 72 °C for 10 min. The PCR procedure for nLSU was as follows: initial denaturation at 94 °C for 1.5 min and a final extension of 72 °C for 10 min. The PCR products were purified and sequenced at Kunming Tsingke Biological Technology Limited Company, Kunming, Yunnan Province, P.R. China. All newly-generated sequences were deposited in NCBI GenBank (Table 1).

The sequences were aligned in MAFFT version 7 (Katoh et al. 2019) using the G-INS-i strategy. The alignment was adjusted manually using AliView version 1.27 (Larsson 2014). Each dataset was aligned separately at first and then the ITS1, 5.8S, ITS2 and nLSU regions were combined with Mesquite version 3.51. The combined dataset was deposited in TreeBASE (submission ID 28564). *Climacocystis borealis* (Fr.) Kotl. and Pouzar and *Diplomitoporus crustulinus* (Bres.) Domański were selected as outgroup (Fig. 1) as inspired by a previous study (Justo et al. 2017).

Maximum parsimony analysis in PAUP\* version 4.0a169 (http://phylosolutions.com/paup-test/) was applied to the combined ITS1+5.8S+ITS2+nLSU dataset. All characters were equally weighted and gaps were treated as missing data. Trees were inferred using the heuristic search option with TBR branch swapping and 1,000 random sequence additions. Max-trees were set to 5,000, branches of zero length were collapsed and all parsimonious trees were saved. Clade robustness was assessed using bootstrap (BT) analysis with 1,000 pseudoreplicates (Felsenstein 1985). Descriptive tree statistics – tree length (TL), composite consistency index (CI), composite retention index (RI), composite rescaled consistency index (RC) and composite homoplasy index (HI) – were calculated for each maximum parsimonious tree generated. The combined dataset was also analysed using Maximum Likelihood (ML) in RAxML-HPC2 through the CIPRES Science Gateway (Miller et al. 2012). Branch support (BS) for the ML analysis was determined by 1,000 bootstrap pseudoreplicates.

6	Sector N			D.C.
Species name	Specimen No.	Gendank a	ccession ino.	References
	777	115	LSU	
Climacocystis borealis	FD-31	KP135308	KP135210	Justo et al. (2017)
Diplomitoporus crustulinus	FD-137	KP135299	KP135211	Justo et al. (2017)
Hyphoderma amoenum	USO 286622	HE577030	-	Telleria et al. (2012)
H. assimile	CBS 125852	MH863808	MH875272	Vu et al. (2019)
H. cremeoalbum	NH 11538	DQ677492	DQ677492	Larsson (2007)
H. crystallinum	CLZhao 9338	MW917161	MW913414	Guan and Zhao (2021a)
	CLZhao 9374	MW917162	MW913415	Guan and Zhao (2021a)
	CL 7bao 1022/	MW/917163	MW/913/16	Guan and Zhao (2021a)
	CLZhao 10224	MW017166	MW/012/17	Guan and Zhao (2021a)
	CLZIIa0 11/23	MW017164	MW012410	Guan and Zhao $(2021a)$
	CLZhao 15841	MW91/165	MW913418	Guan and Zhao (2021a)
	CLZhao 18459	MW917166	MW913419	Guan and Zhao (2021a)
H. definitum	GEL 2898	—	AJ406509	Yurchenko and Wu (2014)
	NH 12266	DQ677493	DQ677493	Larsson (2007)
H. fissuratum	CLZhao 6731	MT791331	MT791335	Ma et al. (2021)
	CLZhao 6726	MT791330	MT791334	Ma et al. (2021)
H. floccosum	CLZhao 17129	MW301683	MW293733	Guan and Zhao (2021b)
	CLZhao 17296	MW301686	MW293736	Guan and Zhao (2021b)
	CL7bao 16/92	MW/301688	MW/29373/	Guan and Zhao (2021b)
	CLZhao 10492	MW/201607	MW/202725	Guan and Zhao (2021b)
11 1.6	CLZIIa0 1/21)	INI W J0108/	INI W 293733	$V_{\rm ext} = 1 $ $V_{\rm ext} = 1 $ $V_{\rm ext} = 1 $
F1. granuujerum	KHL 12561	JIN/10545	JIN/10545	furchenko and Wu (2014)
H. incrustatum	KHL 6685		AY586668	Yurchenko and Wu (2014)
H. litschaueri	NH 7603	DQ677496	DQ677496	Larsson (2007)
	FP-101740-Sp	KP135295	KP135219	Floudas and Hibbett (2015)
H. macaronesicum	MA:Fungi:16099	HE577027	—	Yurchenko and Wu (2014)
	TFC:Mic.15981	HE577028	_	Yurchenko and Wu (2014)
H. medioburiense	NH 10950	DQ677497	DQ677497	Larsson (2007)
H. membranaceum	CLZhao 5844	MW917167	MW913420	Guan and Zhao (2021a)
	CI 7hao 6971	MW917168	MW/913421	Guan and Zhao (2021a)
H microporaides	CLZhao 6857	MW/917169	MW/913/22	Guan and Zhao (2021a)
11. microporotaes	CLZHao 0857	MW017170	MW012422	Guan and Zhao (2021a)
11		MW91/1/0	WIW919422	Guan and Zhao (2021a)
H. moniliforme	Wu 0211–42	KC928282		Yurchenko and Wu (2015)
	Wu 0211–46	KC928284	KC928285	Yurchenko and Wu (2015)
H. mopanshanense	CLZhao 6498	MT791329	MT791333	Ma et al. (2021)
	CLZhao 6493	MT791328	MT791332	Ma et al. (2021)
H. nemorale	TNM F3931	KJ885183	KJ885184	Yurchenko and Wu (2015)
	Wu 9508–14	KC928280	KC928281	Yurchenko and Wu (2015)
H. nudicephalum	Wu 9307–29	AJ534269	_	Nilsson et al. (2003)
I	Wu 9508-225	AI534268	_	Nilsson et al. (2003)
H aptusiforme	KHI 1464	IN572909	_	Yurchenko and Wu (2014)
11. 001431j07me	KHL 11105	INI572010		Yurshenko and Wu (2014)
77 I.	KIIL 1110)	JIN )/2910	AVE 0.((70)	V l l W (2014)
H. obtusum	JS 1/804		AY5866/0	Yurchenko and Wu (2014)
H. occidentale	KHL 8469	—	AY5866/4	Yurchenko and Wu (2014)
	KHL 8477	DQ677499	DQ677499	Larsson (2007)
H. paramacaronesicum	MA:Fungi:87736	KC984399	-	Martín et al. (2018)
	MA:Fungi:87737	KC984405	—	Martín et al. (2018)
H. pinicola	Wu 0108–32	KJ885181	KJ885182	Yurchenko and Wu (2014)
1	Wu 0108–36	KC928278	KC928279	Yurchenko and Wu (2014)
H. prosopidis	E09/58-9	HE577029	_	Yurchenko and Wu (2015)
H puerense	CI 7hao 9476*	MW/443045	_	Present study
II. puctose	CLZhao 9583	MW/4/30/46	MW/4/43051	Present study
11	NUL 105 45	101 10 11 10 10	AVE0((72)	V 1 1 1 W (2014)
H. roseocremeum	NH 10545		AY5866/2	Yurchenko and Wu (2014)
H. setigerum	FCUG 1200	AJ5342/3	_	Nilsson et al. (2003)
H. setigerum	FCUG 1688	AJ534272	—	Nilsson et al. (2003)
H. sinense	CLZhao 7963	MW301679	MW293730	Guan and Zhao (2021b)
	CLZhao 17811	MW301682	MW293732	Guan and Zhao (2021b)
	CLZhao 7981	MW301680	MW293731	Guan and Zhao (2021b)
Hyphoderma sp.	KUC20121102-21	KJ668522	_	Unpublished
*	KUC11052	KI714002	_	Jang et al. (2015)
	Wu 0311_25	KR868735	_	Unpublished
	Wu 0310 6	KR868736	_	Unpublished
	WL 0010-0	KD0/0727	_	Uppublished
	WU U8U8-8/	NR008/3/	_	Unpublished
<b>TT</b> 1 .	GEL3689	DQ34032/	—	Unpublished
H. subsetigerum	Wu 9304–18	AJ534277	_	Nilsson et al. (2003)
	Wu 9202–15	AJ534278	-	Nilsson et al. (2003)

Table 1. List of species, specimens and GenBank accession numbers of sequences used in this study.

Species name	Specimen No.	GenBank accession No.		References
		ITS	LSU	
H. subsetigerum	HHB11620	GQ409521	_	Yurchenko and Wu (2014)
	CFMR MJL1536	GQ409522	_	Yurchenko and Wu (2014)
H. tenuissimum	CLZhao 6930	MW443047	MW443052	Present study
	CLZhao 7003	MW443048	MW443053	Present study
	CLZhao 7221*	MW443049	MW443054	Present study
	CLZhao 16210	MW443050	MW443055	Present study
H. transiens	NH 12304	DQ677504	DQ677504	Larsson (2007)
H. variolosum	CBS 734.91	MH862320	MH873992	Vu et al. (2019)
	CBS 735.91	MH862321	MH873993	Vu et al. (2019)
Hypochnicium erikssonii	NH 9635	_	DQ677508	Larsson (2007)
H. geogenium	NH 10910	_	DQ677509	Larsson (2007)
	MA-Fungi 48308	FN552534	JN939576	Telleria et al. (2010)
H. michelii	MA-Fungi 79155	NR119742	NG060635	Telleria et al. (2010)
H. punctulatum	FP101698sp	KY948827	KY948860	Justo et al. (2017)
H. sphaerosporum	RLG15138sp	KY948803	KY948861	Justo et al. (2017)
H. wakefieldiae	MA-Fungi 7675	FN552531	JN939577	Telleria et al. (2010)
Physisporinus subcrocatus	Dai 15917	KY131870	KY131926	Wu et al. (2017)
P. subcrocatus	Dai 12800	KY131869	KY131925	Wu et al. (2017)
P. tibeticus	Cui 9588	KY131873	KY131929	Wu et al. (2017)
	Cui 9518	KY131872	KY131928	Wu et al. (2017)
Rigidoporus eminens	Dai 17200	MT279690	MT279911	Wu et al. (2017)
R. undatus	Miettinen-13591	KY948731	KY948870	Justo et al. (2017)

New species is shown in bold; \* type material.

MrModeltest 2.3 (Nylander 2004) was used to determine the best-fit evolution model for each dataset (ITS1+5.8S+ITS2+nLSU) for Bayesian Inference (BI). BI was calculated with MrBayes version 3.2.7a (Ronquist et al. 2012). Four Markov chains were run for two runs from random starting trees for 3 million generations (Fig. 1). The first 25% of all generations was discarded as burn-in. A majority rule consensus tree was computed from the remaining trees. Branches were considered as significantly supported if they received a maximum likelihood bootstrap support value (BS) of > 70%, a maximum parsimony bootstrap support value (BT) of > 70% or a Bayesian posterior probability (BPP) of > 0.95.

#### Results

#### Molecular phylogeny

The ITS1+5.8S+ITS2+nLSU dataset comprised sequences from 86 fungal specimens representing 46 taxa. The dataset had an aligned length of 2,034 characters, of which 1,360 characters were constant, 131 were variable and parsimony-uninformative and 543 (35%) were parsimony-informative. Maximum parsimony analysis yielded 108 equally parsimonious trees (TL = 3,317, CI = 0.3361, HI = 0.6946, RI = 0.7051 and RC = 0.2370). The best model of nucleotide evolution for the ITS1+5.8S+ITS2+nLSU dataset estimated and applied in the Bayesian analysis was found to be GTR+I+G. Bayesian analysis and ML analysis resulted in a similar topology as in the MP analysis. The Bayesian analysis had an average standard deviation of split frequencies = 0.008952 (BI) and the effective sample size (ESS) across the two runs is double the average ESS (avg. ESS) = 1,771. The Bayesian tree is shown here (Fig. 1).



**Figure 1.** Maximum parsimony strict consensus tree illustrating the phylogeny of the two new species and related species in *Hyphoderma*, based on ITS1+5.8S+ITS2+nLSU sequences. Branches are labelled with maximum likelihood bootstrap values > 70%, parsimony bootstrap values > 50% and Bayesian posterior probabilities > 0.95, respectively.

The phylogram inferred from ITS1+5.8S+ITS2+nLSU sequences (Fig. 1) highlights the two undescribed species in *Hyphoderma*; *H. puerense* as a sister to *H. moniliforme* and *H. tenuissimum* that forms an independent monophyletic lineage (100% parsimony bootstrap support, 100% likelihood bootstrap support and 1.00 BPP).

## Taxonomy

Hyphoderma puerense C.L. Zhao & Q.X. Guan, sp. nov.

MycoBank No: 838411 Figs 2, 3

**Holotype.** China. Yunnan Province, Puer, Jingdong County, Huilianghe Village, GPS co-ordinates 24°04'45"N, 100°56'32"E, altitude 1246 m a.s.l., on fallen angiosperm branch, leg. C.L. Zhao, 4 January 2019, CLZhao 9476 (SWFC).

Etymology. puerense (Lat.): referring to the locality (Puer) of the specimens.



Figure 2. Basidiomata of Hyphoderma puerense (holotype). Scale bars: 2 cm (a); 1 mm (b).



**Figure 3.** Microscopic structures of *Hyphoderma puerense* (holotype) **a** basidiospores **b** basidia and basidioles **c** cystidia **d** a section of hymenium. Scale bars:  $5 \mu m$  (**a**);  $10 \mu m$  (**b–d**).

**Description.** Basidioma annual, resupinate, adnate, byssoid, without odour and taste when fresh, up to 15 cm long, 3 cm wide, 100–260  $\mu$ m thick. Hymenial surface smooth to floccose, cream when fresh, cream to slightly buff on drying. Margin sterile, thinning out, narrow, cream.

Hyphal system monomitic, generative hyphae with clamps, colourless, thickwalled, frequently branched, interwoven,  $2.5-4.5 \mu m$  in diameter; IKI-, CB-; tissues unchanged in KOH; subhymenial hyphae densely covered by crystals.

Cystidia tubular, encrusted with small crystals,  $25-97 \times 5.5-9.5 \mu m$ .

Basidia clavate to subcylindrical, slightly constricted in the middle to somewhat sinuous, with 4 sterigmata and a basal clamp,  $20-30 \times 4.5-6 \mu m$ .

Basidiospores ellipsoid, colourless, thin-walled, smooth, IKI-, CB-,  $(5.5-)6-7.5(-8) \times 3-4.5(-5) \mu$ m, L = 6.53  $\mu$ m, W = 3.71  $\mu$ m, Q = 1.73-1.79 (n = 60/2).

Habitat and ecology. Climate of the sample collection site is subtropical monsoon climate area, the forest type is evergreen angiosperm forest and samples were collected on fallen angiosperm branches.

Additional specimens examined. China. Yunnan Province, Puer, Jingdong County, Huilianghe Village, GPS co-ordinates 24°04'45"N, 100°56'32"E, altitude 1246 m a.s.l., on fallen angiosperm branch, leg. C.L. Zhao, 5 January 2019, CLZhao 9583 (SWFC).

## Hyphoderma tenuissimum C.L. Zhao & Q.X. Guan, sp. nov.

MycoBank No: 838412 Figs 4, 5

**Holotype.** China. Yunnan Province, Chuxiong, Zixishan Forestry Park, GPS co-ordinates 25°01'26"N, 101°24'37"E, altitude 2313 m a.s.l., on fallen angiosperm branch, leg. C.L. Zhao, 1 July 2018, CLZhao 7221 (SWFC).

Etymology. tenuissimum (Lat.): referring to the thin basidiomata.



Figure 4. Basidiomata of Hyphoderma tenuissimum (holotype). Scale bars: 2 cm (a); 1 mm (b).



**Figure 5.** Microscopic structures of *Hyphoderma tenuissimum* (holotype) **a** basidiospores **b** basidia and basidioles **c** cystidia **d** a section of hymenium. Scale bars: 10  $\mu$ m (**a**–**d**).

**Description.** Basidioma annual, resupinate, adnate, membranaceous when fresh, hard membranaceous upon drying, up to 20 cm long, 3 cm wide,  $30-100 \mu m$  thick. Hymenial surface tuberculate to minutely-grandinioid, slightly buff when fresh, buff upon drying, cracking. Margin sterile, slightly buff, 1 mm wide.

Hyphal system monomitic, generative hyphae with clamps, colourless, thick-walled, frequently branched, interwoven, 3–5 µm in diameter, IKI-, CB-; tissues unchanged in KOH.

Cystidia large, cylindrical, with 4–12 clamped septa, with abundant encrustations, 50–220  $\times$  6.5–13  $\mu m.$ 

Basidia clavate to subcylindrical, constricted, somewhat sinuous, with 4 sterigmata and a basal clamp connection,  $17-31 \times 4.5-8 \mu m$ .

Basidiospores cylindrical, colourless, thin-walled, smooth, with oil drops inside, IKI–, CB–, 7–10.5(–11) × 3–4.5(–5)  $\mu$ m, L = 8.75  $\mu$ m, W = 4.15  $\mu$ m, Q = 2.02–2.18 (n = 120/4).

Habitat and ecology. Climate of the sample collection site is subtropical monsoon climate area, the forest type is evergreen angiosperm forest and samples were collected on fallen angiosperm branches.

Additional specimens examined. China. Yunnan Province, Chuxiong, Zixishan National Forestry Park, GPS co-ordinates 25°01'26"N, 101°24'37"E, altitude 2263 m a.s.l., on fallen angiosperm branch, leg. C.L. Zhao, 1 July 2018, CLZhao 6930, CLZhao 7003 (SWFC); Wenshan, Pingba Town, Wenshan National Nature Reserve, GPS co-ordinates 23°18'19"N, 104°42'47"E, altitude 1976 m a.s.l., on fallen angiosperm branch, leg. C.L. Zhao, 25 July 2019, CLZhao 16210 (SWFC).

## Discussion

In the present study, two new species, *Hyphoderma puerense* and *H. tenuissimum* are described, based on phylogenetic analyses and morphological characters.

Phylogenetically, the two new taxa were found to belong to *Hyphoderma*, in which *H. puerense* forms a sister species to *H. moniliforme* and *H. tenuissimum* forms an independent monophyletic lineage (100% BS, 100% BP and 1.00 BPP).

Morphologically, *Hyphoderma puerense* is similar to *H. obtusiforme* J. Erikss. & Å. Strid and *H. obtusum* in having a smooth hymenium, non-septate cylindrical cystidia and ellipsoid basidiospores. However, *H. obtusiforme* differs from *H. puerense* by both larger basidia ( $30-40 \times 8-9 \mu m$ ) and basidiospores ( $10-14 \times 5-7 \mu m$ ; Eriksson and Ryvarden 1975). *Hyphoderma obtusum* also differs from *H. puerense* by larger basidia ( $30-35 \times 6-8 \mu m$ ) and basidiospores ( $8-9 \times 5-6.5 \mu m$ ; Eriksson 1958). *Hyphoderma puerense* is similar to *H. roseocremeum* (Bres.) Donk in having smooth hymenium and non-septate cylindrical cystidia. However, *Hyphoderma roseocremeum* differs through the presence of larger basidiospores ( $8-12 \times 3-4 \mu m$ ; Bernicchia and Gorjón 2010).

Morphologically, *Hyphoderma tenuissimum* is similar to *H. floccosum* C.L. Zhao & Q.X. Guan, *H. mopanshanense*, *H. nudicephalum* Gilb. & M. Blackw., *H. pinicola*, *H. setigerum* and *H. subsetigerum* Sheng H. Wu in having septocystidia and cylindrical basidiospores. However, *Hyphoderma floccosum* differs from *H. tenuissimum* by having a floccose hymenial surface and tubular cystidia (Guan and Zhao 2021b); *H. mopanshanense* is separated from *H. tenuissimum* by having porulose to pilose hymenial surface and smaller basidia (15–18.5 × 3–4.5 µm; Ma et al. 2021); *H. nu*-

*dicephalum* differs from *H. tenuissimum* in the nature of the septocystidial apex (lacking encrustation; swollen up to 14  $\mu$ m; Gilbertson and Blackwell 1988); *H. pinicola* is separated from *H. tenuissimum* by having basidia with two sterigmata and larger basidiospores (13–16 × 4–4.5  $\mu$ m; Yurchenko and Wu 2014b); *H. setigerum* differs by having a combination of thin basidiomata with very long septocystidia (Bernicchia and Gorjón 2010); and *H. subsetigerum* differs from *H. tenuissimum* by having narrower basidia (20–30 × 4.5–5.5  $\mu$ m) and smaller basidiospores (6–8 × 2.8–3.2  $\mu$ m; Wu 1997).

Nilsson et al. (2003) highlighted the phylogeography of *Hyphoderma setigerum* (Basidiomycota) in the Northern Hemisphere in a study based on molecular analysis, morphological studies and crossing tests. Nine preliminary taxa were shown to exist inside the *H. setigerum* complex; in the present study, *H. tenuissimum* belongs to the *H. setigerum* complex, based on the morphological character of long septocystidia and phylogenetic evidence. A previous study indicated the importance of vicariance in the evolution of this species complex (Nilsson et al. 2003) and our study shows that the specimens of *H. tenuissimum* are collected in Zixishan National Forestry Park (GPS co-ordinates 25°01'26"N, 101°24'37"E), Chuxiong, Yunnan Province, China, which is distinct from *H. setigerum* s. str. (Norway: Oppland and Finland: Pohjois-Häme). The present samples of *H. subsetigerum* and *H. nudicephalum* were collected in Yunnan Province, China, but neither of these taxa groups together closely with *H. tenuissimum* (Fig. 1).

In the current phylogenetic tree, two partially annotated GenBank sequences (KJ668522 and KJ714002) of *Hyphoderma* sp. (South Korea) cluster closely with four sequences of the new species *Hyphoderma tenuissimum*, although whether they really belong to this species remains to be assessed. It is certainly conceivable that they do, which would mean that *Hyphoderma tenuissimum* has been collected and sequenced at least six times in Asia. Regarding the new taxon *H. puerense* (Fig. 1), four partially annotated GenBank sequences (KR868735, KR868736, KR868737 and DQ340327) form a reasonably well-supported clade together with our two specimens of *H. puerense*. We interpret this to mean that all six taxa represent *H. puerense*. All of the samples in this clade are from Asia, which supports the point of the importance of vicariance in the evolution in this genus.

### Key to 30 accepted species of Hyphoderma in China

1	Cystidia absent	2
_	Cystidia present	5
2	Hymenial surface grandinioid	H. acystidiatum
_	Hymenial surface smooth	
3	Basidiospores > 10.5 µm in length	H. densum
_	Basidiospores < 10.5 µm in length	4
4	Hymenophore cracked; basidiospores > 8.5 µm in length	H. fissuratum
_	Hymenophore uncracked; basidiospores < 8.5 µm in length	H. sibiricum
5	Hymenophore smooth	6
_	Hymenophore tuberculate, porulose, grandinioid or odontoic	l15

6	Two types of cystidia present	7
_	One type of cystidia present	8
7	Moniliform cystidia absent	.H. microcystidium
_	Moniliform cystidia present	H. sinense
8	Hymenophore uncracked	9
_	Hymenophore cracked	
9	Basidiospores > 11 µm in length	H. definitum
_	Basidiospores < 11 µm in length	
10	Basidiospores > 8.5 µm in length	H. microporoides
_	Basidiospores < 8.5 µm in length	H. puerense
11	Cystidia moniliform	
_	Cystidia cylindrical	
12	Basidiospores > 9 μm in length	H. litschaueri
_	Basidiospores < 9 μm in length	H. moniliforme
13	Basidiospores ellipsoid < 10 µm in length	H. rimulosum
_	Basidiospores cylindrical > 10 µm in length	14
14	Basidiospores > 12 µm in length	H. cremeum
_	Basidiospores < 12 µm in length	H. subclavatum
15	Hymenophore odontoid or grandinioid	16
_	Hymenophore tuberculate, porulose	19
16	Hymenophore odontoid	
_	Hymenophore grandinioid	
17	Basidiospores < 4.5 µm in width	H. transiens
_	Basidiospores > 4.5 µm in width	H. formosanum
18	Basidiospores larger 7–10.5 × 3–4.5 $\mu$ m	H. tenuissimum
_	Basidiospores smaller $6-8 \times 2.8-3.2 \mu\text{m}$	H. subsetigerum
19	Cystidia of two types	20
_	Cystidia of one type	23
20	Septate cystidia absent	21
_	Septate cystidia present	
21	Basidiospores $< 4  \mu m$ in width	H. variolosum
_	Basidiospores > 4 $\mu$ m in width	H. crystallinum
22	Basidia 2-sterigmata, basidiospores > 13 µm in length	
_	Basidia 4-sterigmata, basidiospores < 13 µm in length	
23	Septate cystidia present	
_	Septate cystidia absent	25
24	Hymenophore porulose to pilose, basidia < 5 µm in width	.H. mopanshanense
_	Hymenophore tuberculate, basidia > 5 µm in width	
25	Hymenophore porulose	H. obtusiforme
_	Hymenophore tuberculate, colliculose	
26	Cystidia < 30 μm in length	H. cremeoalbum
_	Cystidia > 30 µm in length	
27	Basidia > 30 µm in length	
_	Basidia < 30 µm in length	29

28	Hymenophore cracking, cystidia < 10 µm in width	H. medioburiense
_	Hymenophore not cracking, cystidia > 10 µm in width	H. clavatum
29	Hymenophore colliculose	H. nemorale
_	Hymenophore tuberculate	H. membranaceum

## Acknowledgements

The research was supported by the Yunnan Fundamental Research Project (Grant No. 202001AS070043) and Science Research Foundation of Yunnan Provincial Department of Education Project (Project No. 2021Y275).

## References

- Baltazar JM, Silveira RMB, Rajchenberg M (2016) Type studies of J. Rick's corticioid homobasidiomycetes (Agaricomycetes, Basidiomycota) housed in the Herbarium Anchieta (PACA). Phytotaxa 255(2): 101–132. https://doi.org/10.11646/phytotaxa.255.2.1
- Bernicchia A, Gorjón SP (2010) Fungi Europaei 12: Corticiaceae s.l. Alassio: Edizioni Candusso.
- Dai YC (2012) Polypore diversity in China with an annotated checklist of Chinese polypores. Mycoscience 53(1): 49–80. https://doi.org/10.1007/s10267-011-0134-3
- Donk MA (1957) Notes on resupinate Hymenomycetes IV. Fungus 27: 1-29.
- Eriksson J (1958) Studies in the Heterobasidiomycetes and Homobasidiomycetes Aphyllophorales of Muddus National Park in North Sweden. Symbolae Botanicae Upsalienses 16(1): 1–172.
- Eriksson J, Ryvarden L (1975) The Corticiaceae of North Europe; Fungiflora: Oslo, Norway, Volume 3, 288–546.
- Felsenstein J (1985) Confidence intervals on phylogenetics: An approach using bootstrap. Evolution 39: 783–791. https://doi.org/10.1111/j.1558-5646.1985.tb00420.x
- Floudas D, Hibbett DS (2015) Revisiting the taxonomy of *Phanerochaete* (Polyporales, Basidiomycota) using a four gene dataset and extensive ITS sampling. Fungal Biology 119(80): 679–719. https://doi.org/10.1016/j.funbio.2015.04.003
- Gilbertson RL, Blackwell M (1988) Some new or unusual corticioid fungi from the Gulf Coast region. Mycotaxon 33: 375–386.
- Guan QX, Zhao CL (2021a) Taxonomy and phylogeny of the wood-inhabiting fungal genus *Hyphoderma* with descriptions of three new species from East Asia. Journal of Fungi 7(4): e308. https://doi.org/10.3390/jof7040308
- Guan QX, Zhao CL (2021b) Two new corticioid species, *Hyphoderma sinense* and *H. floccosum* (Hyphodermataceae, Polyporales), from southern China. Mycosystema 40(03): 447–461. https://doi.org/10.13346/j.mycosystema.200382
- James TY, Stajich JE, Hittinger CT, Rokas A (2020) Toward a fully resolved fungal tree of life. Annual Review of Microbiology 74(1): 291–313. https://doi.org/10.1146/annurevmicro-022020-051835
- Jang Y, Jang S, Min M, Hong JH, Lee H, Lee H, Lim YW, Kim JJ (2015) Comparison of the diversity of basidiomycetes from dead wood of the Manchurian fir (*Abies holophylla*) as

evaluated by fruiting body collection, mycelial isolation, and 454 sequencing. Microbial Ecology 70(3): 634–645. https://doi.org/10.1007/s00248-015-0616-5

- Justo A, Miettinen O, Floudas D, Ortiz-Santana B, Sjökvist E, Linder D, Nakasone K, Niemelä T, Larsson K, Ryvarden L, Hibbetta DS (2017) A revised family-level classification of the Polyporales (Basidiomycota). Fungal Biology 121(9): 798–824. https://doi.org/10.1016/j. funbio.2017.05.010
- Katoh K, Rozewicki J, Yamada KD (2019) MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. Briefings in Bioinformatics 20(4): 1160–1166. https://doi.org/10.1093/bib/bbx108
- Larsson A (2014) AliView: a fast and lightweight alignment viewer and editor for large data sets. Bioinformatics 30(22): 3276–3278. http://dx.doi.org/10.1093/bioinformatics/btu531
- Larsson KH (2007) Re-thinking the classification of corticioid fungi. Mycological Research 111(9): 1040–1063. https://doi.org/10.1016/j.mycres.2007.08.001
- Ma X, Huang RX, Zhang Y, Zhao CL (2021) *Hyphoderma fissuratum* and *H. mopanshanense* spp. nov. (Polyporales) from southern China. Mycoscience 62(1): 36–41. https://doi. org/10.47371/Mycosci.2020.08.004
- Martín MP, Zhang LF, Fernández-López J, Dueñas M, Rodríguez-Armas JL, Beltrán-Tejera E, Telleria MT (2018) *Hyphoderma paramacaronesicum* sp. nov. (Meruliaceae, Polyporales, Basidiomycota), a cryptic lineage to *H. macaronesicum*. Fungal Systematics and Evolution 2: 57–68. https://doi.org/10.3114/fuse.2018.02.05
- Miller MA, Pfeiffer W, Schwartz T (2012) The CIPRES Science Gateway: enabling high-impact science for phylogenetics researchers with limited resources. Association for Computing Machinery 39: 1–8. https://doi.org/10.1145/2335755.2335836
- Nakasone KK (2008) Type studies of corticioid Hymenomycetes described by Bresadola. Cryptogamie Mycologie 29(3): 231–257. https://doi.org/10.1002/yea.1629
- Nilsson RH, Hallenberg N, Nordén B, Maekawa N, Wu SH (2003) Phylogeography of *Hyphoderma setigerum* (Basidiomycota) in the Northern Hemisphere. Mycological Research 107(6): 645–652. https://doi.org/10.1017/S0953756203007925
- Nylander JAA (2004) MrModeltest v.2. Program Distributed by the Author. Evolutionary Biology Centre, Uppsala University, Uppsala.
- Petersen JH (1996) The Danish Mycological Society's colour-chart. Foreningen til Svampekundskabens Fremme, Greve.
- Rehner SA, Samuels GJ (1994) Taxonomy and phylogeny of *Gliocladium* analysed from nuclear large subunit ribosomal DNA sequences. Mycological Research 98(6): 625–634. https:// doi.org/10.1016/S0953-7562(09)80409-7
- Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Hohna S, Larget B, Liu L, Suchard MA, Huelsenbeck JP (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. Systematic Biology 61(3): 539–542. https://doi.org/10.1093/sysbio/sys029
- Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios AM, Thu PQ, Suija A, Smith ME, Sharp C, Saluveer E, Saitta A, Rosas M, Riit T, Ratkowsky D, Pritsch K, Põldmaa K, Piepenbring M, Phosri C, Peterson M, Parts K, Pärtel K, Otsing E, Nouhra E, Njouonkou AL, Nilsson RH, Morgado LN, Mayor J, May TW, Majuakim L, Lodge DJ, Lee SS, Larsson K-H, Kohout P, Hosaka K, Hiiesalu I, Henkel

TW, Harend H, Guo L-d, Greslebin A, Grelet G, Geml J, Gates G, Dunstan W, Dunk C, Drenkhan R, Dearnaley J, De Kesel A, Dang T, Chen X, Buegger F, Brearley FQ, Bonito G, Anslan S, Abelland S, Abarenkov K (2014) Global diversity and geography of soil fungi. Science 346: 1256688. https://doi.org/10.1126/science.1256688

- Telleria MT, Dueñas M, Beltrán-Tejera E, Rodríguez-Armas JL, Martín MP (2012) A new species of *Hyphoderma* (Meruliaceae, Polyporales) and its discrimination from closely related taxa. Mycologia 104(5): 1121–1132. https://doi.org/10.3852/11-344
- Telleria MT, Duenas M, Melo I, Hallenberg N, Martin MP (2010) A re-evaluation of *Hypochnicium* (Polyporales) based on morphological and molecular characters. Mycologia 102(6): 1426–1436. https://doi.org/10.3852/09-242
- Vilgalys R, Hester M (1990) Rapid genetic identification and mapping of enzymatically amplified ribosomal DNA from several *Cryptococcus* species. Journal of Bacteriology 172(8): 4238–4246.
- Vu D, Groenewald M, Vries M, Gehrmann T, Stielow B, Eberhardt U, Al-Hatmi A, Groenewald JZ, Cardinali G, Houbraken J, Boekhout T, Crous PW, Robert V, Verkley GJM (2019) Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom Fungi and reveals thresholds for fungal species and higher taxon delimitation. Studies in Mycology 92: 135–154. https://doi.org/10.1016/j.simyco.2018.05.001
- Wallroth CFW (1833) Flora Cryptogamica Germaniae 2 Algas et fungos. Norimbergae [Nüremberg], Schragius [J.L. Schrag], 923 pp.
- White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In: Innis MA, Gefand DH, Sninsky JJ, White JT (Eds) PCR Protocols: A Guide to Methods and Applications. Academic Press, San Diego, 18: 315–322. https://doi.org/10.1016/B978-0-12-372180-8.50042-1
- Wu F, Chen JJ, Ji XH, Vlasák J, Dai YC (2017) Phylogeny and diversity of the morphologically similar polypore genera *Rigidoporus*, *Physisporinus*, *Oxyporus* and *Leucophellinus*. Mycologia 109(5): 749–765. doi:10.1080/00275514.2017.1405215
- Wu SH (1997) New species of *Hyphoderma* from Taiwan. Mycologia 89: 132–140. https://doi. org/10.1080/00275514.1997.12026764
- Wu SH, Nilsson HR, Chen CT, Yu SY, Hallenberg N (2010) The white-rotting genus *Phanerochaete* is polyphyletic and distributed throughout the phlebioid clade of the Polyporales (Basidiomycota). Fungal Diversity 42(1): 107–118. https://doi.org/10.1007/s13225-010-0031-7
- Yurchenko E, Wu SH (2014) Hyphoderma pinicola sp. nov. of H. setigerum complex (Basidiomycota) from Yunnan, China. Botanical Studies 55: 71–78. https://doi.org/10.1186/ s40529-014-0071-5
- Yurchenko E, Wu SH (2015) Hyphoderma moniliforme and H. nemorale (Basidiomycota) newly recorded from China. Mycosphere 6(1): 113–121. https://doi.org/10.5943/mycosphere/6/1/11
- Zhao CL, Wu ZQ (2017) Ceriporiopsis kunmingensis sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. Mycological Progress 16(1): 93–100. https://doi.org/10.1007/s11557-016-1259-8