

Volume 14 - Issue 2 | February 2022

mdpi.com/journal/diversity ISSN 1424-2818

MDPI



# Three New Species of Absidia (Mucoromycota) from China Based on Phylogeny, Morphology and Physiology

by 😵 Heng Zhao, 😢 Yong Nie, 😵 Tongkai Zong, 😵 Yucheng Dai and 🤔 Xiaoyong Liu Diversity 2022, 14(2), 132; https://doi.org/10.3390/d14020132 - 13 Feb 2022 Viewed by 322

Abstract Species of Absidia are distributed widely in the environment, while their diversity is insufficiently studied. Three new species, A. frigida, A. gemella and A. longissima, are proposed herein from Xinjiang and Yunnan in China based on phylogenetic, morphological and physiological evidence. [...] Read more.

(This article belongs to the Special Issue Diversity and Evolution of Fungi)

```
► Show Figures
```

#### Open Access Article

# Diversity of Wood-Decaying Fungi in Wuliangshan Area, Yunnan Province, P.R. China

by 😫 Xiao He and 😩 Chang-Lin Zhao

Diversity 2022, 14(2), 131; https://doi.org/10.3390/d14020131 - 11 Feb 2022. Viewed by 249

Abstract Five surveys were carried out in the Wullangshan area, Yunnan Province, P.R. China, based on a combination of morphological features and molecular evidence. Around 2454 specimens of wood-decaying fungi were collected. The paper summarizes the obtained results on the wood-decaying fungi of this [...] Read more.

(This article belongs to the Special Issue Diversity and Evolution of Fungi)

► Show Figures

Open Access Article

# Human-Wildlife Conflict at a Suburban–Wildlands Interface: Effects of Short- and Long-Distance Translocations on Red Diamond Rattlesnake (*Crotalus ruber*) Activity and Survival

by 🔁 Aaron G. Corbit and 🚭 William K. Hayes Diversity 2022, 14(2), 130, https://doi.org/10.3390/d14020130 - 11 Feb 2022 Viewed by 402

Abstract The mitigation of human-rattlesnake conflicts often involves euthanizing or translocating the offending rattlesnake. Although translocation is generally considered more humane, especially by the general public, it may negatively impact the translocated individual and may not be effective if that individual returns to areas [...] Read more. (This article belongs to the Special Issue Advances in the Conservation and Ecology of Rattlesnakes)

▶ Show Figures

## Open Access Article



= + m

4 3

# Recruitment Patterns and Potential Climate Change Impacts on Three Florida Hylids with Different Life Histories

by (1) Sky T Button, (2) Cathryn H. Greenberg and (2) James D. Austin Diversity 2022, 14(2), 129. https://doi.org/10.3390/d14020128 - 10 Feb 2022 Viewed by 297

Abstract Altered weather patterns associated with climate change are likely to adversely affect amphibian recruitment, especially for species dependent on ephemeral, geographically isolated wetlands for breeding. Future changes in temperature and rainfall patterns could affect hydroregimes (periodicity, depth, duration, and timing of water in [\_\_] Read more.

(This article belongs to the Special Issue Amphibian Ecology in Geographically Isolated Wetlands)

► Show Figures





# Article Diversity of Wood-Decaying Fungi in Wuliangshan Area, Yunnan Province, P.R. China

Xiao He<sup>1,2</sup> and Chang-Lin Zhao<sup>1,2,3,4,\*</sup>

- Key Laboratory for Forest Resources Conservation and Utilization in the Southwest Mountains of China, Ministry of Education, Southwest Forestry University, Kunming 650224, China; fungixiaohe@163.com
- <sup>2</sup> College of Biodiversity Conservation, Southwest Forestry University, Kunming 650224, China
- <sup>3</sup> Yunnan Key Laboratory for Fungal Diversity and Green Development, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650201, China
- <sup>4</sup> School of Life Sciences, Tsinghua University, Beijing 100084, China
- Correspondence: fungi@swfu.edu.cn

**Abstract:** Five surveys were carried out in the Wuliangshan area, Yunnan Province, P.R. China, based on a combination of morphological features and molecular evidence. Around 2454 specimens of wood-decaying fungi were collected. The paper summarizes the obtained results on the wood-decaying fungi of this area, consisting in 95 species distributed in 59 genera, 23 families and 9 orders. Their hosts and substrates were also identified. A checklist of wood-decaying fungi is given. Sequences of the ITS nrRNA gene region of the studied specimens were generated and phylogenetic analysis was performed with maximum likelihood, maximum parsimony and Bayesian inference methods. The present list of wood-decaying fungi enriches the knowledge of fungal diversity worldwide and supplies the basic data for future applications.

Keywords: Basidiomycota; biological resources; diversity; mycota; Wuliangshan area



Citation: He, X.; Zhao, C.-L. Diversity of Wood-Decaying Fungi in Wuliangshan Area, Yunnan Province, P.R. China. *Diversity* **2022**, *14*, 131. https://doi.org/10.3390/d14020131

Academic Editors: Ipek Kurtboke and Michael Wink

Received: 28 January 2022 Accepted: 9 February 2022 Published: 11 February 2022

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

## 1. Introduction

Fungi are a diverse, monophyletic group of eukaryotes which have immense ecological and economic impacts. These organisms play an important role in ecosystems as diverse as soil, leaves, trees, wood, and hidden layers within their substrate [1]. Many fungi are microscopic or have cryptic life cycles that make detection difficult, and approximately 140 thousand species of fungi have been described, but the potential biodiversity of the group is likely to be in millions of species [2–4]. The diversity for flora of seed plants in the Yunnan Province is higher than other areas in China, and the endemic species of woody plants are rich, in which both supply good substrates for wood-decaying fungi. Wood-decaying fungi include most basidiomycetes and ascomycetes that grow on various kinds of wood, such as living trees, dead standing trees, fallen trunks, fallen branches and stumps, which can be used for industrial, medicinal, edible and economic value [5–10], and display a considerable ability to transform or degrade different environmental contaminants on their extensive organic compound degradation abilities [1].

The Wuliangshan area is located in the south-central part of the Yunnan Province, China, including the Wuliangshan National Nature Reserve and its surrounding areas, named for its location in the Wuliangshan Mountains. The geographical location is between 100°18′–101°13′ E and 23°12′–24°55′ N [11,12] with an altitude of 2000–3300 m and average annual precipitation more than 1500 mm [11,13]. The plant resources are rich in the Wuliangshan area, with 209 families, 1039 genera and 2574 species of seed plants [11,14]. The mid-mountain wet, semi-wet evergreen broad-leaved forest and monsoon evergreen broad-leaved forest are the main forest communities as the dominant species, which increase the fungal diversity of this area [11,12,15].

The wood-decaying fungi are mostly reported in northwest Yunnan Province, and few polypore and corticioid fungi have been reported there so far. According to the modern taxonomy, wood-decaying fungi mainly belong to ten orders of Agaricomycetes, viz., Agaricales, Auriculariales, Cantharellales, Corticiales, Gloeophyllales, Hymenochaetales, Polyporales, Russulales, Thelephorales and Trechisporales [6,7,16]. Therefore, the current wood-decaying fungi catalogues include poroid and corticioid hymenophores. In the present study, five field trips were carried out in the Wuliangshan area, and about 95 species which are distributed in 59 genera, 23 families, and 9 orders were identified from these materials. The aim of this work was to provide an updated checklist and mapping of wood-decaying fungi in the area, as well as to enrich the knowledge of the fungal diversity in this area.

### 2. Materials and Methods

#### 2.1. Sample Collection and Herbarium Specimen Preparation

Fresh fruiting bodies of basidiomycetous macrofungi growing on angiosperm stumps, trunks and branches were collected by the systematically surveyed selected cells in the macroarea through the woods for 5 sampling points (approximately the four corners of the square and center) within  $1 \times 1 \text{ km}^2$  2–3 times every year in the Wuliangshan area of Yunnan Province from October 2017 to January 2019. The fruiting bodies require three basidiomata at least and each individual is up to 3 cm long, 2 cm wide, in which the smallest fruiting structure collected have to include the hymenophore. The samples were photographed in situ and their fresh macroscopic details were recorded. Photographs were recorded by a Jianeng 80D camera. All photos were focus stacked and merged using Helicon Focus software. Macroscopic details were recorded and transported to a field station where the fruit bodies were dried on an electronic food dryer (Fsfruit, Foshan, China) at 45 °C for 48 h. Once dried, the specimens were sealed in envelope and Ziplock plastic bags and labeled. The dried specimens were deposited in the herbarium of Southwest Forestry University (SWFC), Kunming, Yunnan Province.

#### 2.2. Morphological Studies

Macromorphological descriptions were based on field notes. Color terms are from Petersen [17]. Micromorphological data were obtained from the dried specimens and observed under a light microscope following Dai [6]. The following abbreviations were used for the micro characteristics descriptions: KOH = 5% potassium hydroxide, CB = Cotton Blue, CB = acyanophilous, CB + cyanophilous, IKI = Melzer's reagent, IKI - = both inamyloid and indextrinoid, L = mean spore length (arithmetic average of 30 spores), W = mean spore width (arithmetic average of 30 spores), Q = variation in the L/W rationes between the specimens studied, n (a/b) = number of spores (a) measured from given number (b) of specimens.

#### 2.3. Molecular Procedures and Phylogenetic Analysis

CTAB rapid plant genome extraction kit-DN14 (Aidlab Biotechnologies Co., Ltd., Beijing, China) was used to obtain genomic DNA from dried specimens according to the manufacturer's instructions. ITS region was amplified with primer pair ITS5 and ITS4 [18]. The PCR procedure for ITS was as follows: initial denaturation at 95 °C for 3 min, followed by 35 cycles at 94 °C for 40 s, 58 °C for 45 s and 72 °C for 1 min, and a final extension of 72 °C for 10 min. The PCR products were purified using a QIAquick PCR purification kit (Qiagen Inc., Valencia, CA, USA) and directly sequenced at Kunming Tsingke Biological Technology Limited Company, Kunming Yunnan Province, P.R. China. All newly generated sequences were deposited at GenBank (Table 1).

Species Name         Species No.         TS         References         Country           Abundingene specifies         China Varanan         China Varanan         China Varanan         China Varanan         China Varanan           Appendent         China Varanan         China Varanan         China Varanan         USA. Reheval         Reheval         USA. Reheval </th <th></th> <th></th> <th>GenBank Accession No.</th> <th></th> <th colspan="3" rowspan="2">Country</th>			GenBank Accession No.		Country		
Abundispense spericiola         CLZbas 10236         MT19978         Present study         Chrins, Yunnam           Agarrias compositiv         MA Fung 80996         NR, L51745         Unpublished         USA, Bethesda           Alteradictos throyasus         CLZbas 04333         MKA Stats         Present study         Chrins, Yunnam           A. crassitus         CLZbas 0475         MK14101         Present study         Chrins, Yunnam           A. crassitus         SFC 3018         MT81058         [19]         Asperitor, Specific Action, Yunnam           A. artirolitis         ULZbas 0440         MK54368         Present study         Chrins, Yunnam           A. artirolitis         ULZbas 0421         MK54368         Present study         Chrins, Yunnam           A. subroses         CLZbas 0472         MK543689         Present study         Chrins, Yunnam           A. subroses         CLZbas 0472         MK543689         Present study         Chrins, Yunnam           A. subroses         CLZbas 0472         MK543689         Present study         Chrins, Yunnam           A. subroses         CLZbas 0472         MK543689         Present study         Chrins, Yunnam           A. subroses         CLZbas 0472         MK543689         Present study         Chrins, Yunnam <t< th=""><th>Species Name</th><th>Specimen No.</th><th>ITS</th><th>References</th></t<>	Species Name	Specimen No.	ITS	References			
A. generaties         Date 13944         KC 415907         In 1715           Agerics competities         MA Enryl (1996)         NR. 151745         Unpublished         USA, Rehesda           A. consisties         CIZbao 4333         MK434341         Present study         China, Yunnan           A. consisties         CIZbao 433         MK434342         Present study         China, Yunnan           A. consisties         SPG 3004         MR31058         [19]         China, Yunnan           A. consisties         SPG 3004         MR31058         [21]         China, Kunnan           A. staffilture         CIZbao 4325         MK4109173         [21]         China, Yunnan           A. staffilture         CIZbao 472         MK543482         Present study         China, Yunnan           A. staffineere         CIZbao 472         MK543489         Present study         China, Yunnan           A. staffineere         CIZbao 472         MK543481         Present study         China, Yunnan           A. staffineere         CIZbao 472         MK543489         Present study         China, Nunnan           A. staffineere         CIZbao 432         MK543481         Present study         China, Nunnan           A. staffineere         CIZbao 431         MK5023146	Abundienorus quarcicola	CI 7hao 10238	MT150078	Procont study	China Yunnan		
Apprices campestris         MA Farg/180796         NR. 151745         Unpublished         USA, Rethesda           Alcrodiscio korpouse         CLZbao 4733         MK343481         Present study         Chrin, Yunnan           A. crossitis         CLZbao 4735         MK404310         Present study         Chrin, Yunnan           A. crossitis         SFC 3004         MT831058         [19]         Chrin, Yunnan           A. stabellinis         CLZbao 4705         MK434881         Present study         Chrin, Yunnan           A. stabellinis         ILZbao 4621         MK434881         Present study         Chrin, Junnan           A. natabilis         JMB 2152         MT565513         [19]         Chrin, Junnan           A. natabilis         CLZbao 4212         MK26352         Present study         Chrin, Junnan           A. dicitatis         CLZbao 4210         ME241891         [21]         Chrin, Suman           A. dicitatis         CLZbao 4210         ME345353	A auercicola	Dai 3084	KC415907	[19]	China Yunnan		
Absorb         CIPE 356.6         ME4878812         Cip         Cip         Netherlands, Upsallalan           A. Joryosuis         CLZhao 433         MEA04310         Present study         China, Yunnan           A. crussitus         SPG 300         MT831088         [19]         Agenin, Esquel           A. stabilitus         CLZhao 4315         MK404310         Present study         China, Yunnan           A. stabilitus         CLZhao 422         ME109953         [21]         China, Sequel           A. mindulis         CLZhao 423         ME30438         Present study         China, Yunnan           A. mindulis         CLZhao 423         ME119955         [21]         China, Feingel           A. mindulis         CLZhao 3822         ME119955         [21]         China, Feingel           A. distinge         SPC 2018014-01         MK992816         [22]         Kora, Secul           Auriculum issiftito         CLZhao 3425         MK243592         Present study         China, Yunnan           A. distiftita         ON1932         ME48359         Present study         China, Yunnan           A. distiftita         ON1932         ME483592         Present study         China, Yunnan           A. distiftita         ON1932         ME483592	Agaricus campetris	MA Fungi 80998	NR 151745	Unpublished	USA Bethesda		
Accense         Classes         Mickassa         Process         Part         Process         Part         Part<	Alguradiagua batruagua	CBS 226 66	ML959912		Nothorlanda Unsalalaan		
A. Dompsone     CLZIAU 4703     MK404130     Present study     Chima, Iumani       A. crossbirs     SPG 3004     MTS1088     Present study     Chima, Engled       A. sizelefinus     CLZao 4405     MK534382     Present study     China, Jeping       A. sizelefinus     CLZao 4405     MK534382     Present study     China, Jeping       A. minehilis     CLZao 4421     MK343488     Present study     China, Jeping       A. minehilis     CLZao 4722     MK534389     Present study     China, Steping       A. subroseus     CLZao 3832     MH114612     Unpublished     China, Yunnan       A. subroseus     He 4814     MH1109055     [21]     China, Beijing       Antical applicita     SEC 20180314-01     MK992816     [22]     Korae, Secol       Antical applicita     SEC 20180314-01     MK992816     [23]     China, Yunnan       A. delicatin     CLZao 3852     MK204844     Present study     China, Yunnan       A. delicatin     CLZao 3852     MK204316     [23]     Thailand, China, Reijing       A. delicatin     MH114102112     KX621146     [24]     China, Yunnan       A. delicatin     MH114102112     KX621146     [25]     USA, Worcester       B. datasta     CLZhou 4310     MK534505     Presen	A hotmoous	CL Zhao 4222	NIF1030012 MV242481	[20] Procent study	China Vunnan		
A. 6199000000     Clauda, 1010000       A. 6199000000     Clauda, 1010000       A. sidelfunes     CLZhao, 4005       M. Sidelfunes     CLZhao, 4005       M. Sidelfunes     CLZhao, 4005       M. sidelfunes     CLZhao, 4021       M. sidelfunes     CLAB, 4021       M. sidelfunes     CLZhao, 4022       M. Sidelfunes     CLAB, 4021       M. sidelfunes     CLAB, 4023       M. sidelfunes     CLAB, 4023       M. sidelfunes     MK543502       M. sidelfunes     CLAB, 4023       M. sidelfunes     CLZhao, 4245       M. sidelfune     CLAB, 4023       M. sidelfunes     CLAB, 4023       M. Sidelfune	A. boiryosus	CLZIIdo 4555	ME 404210	Procent study	China, Tunnan		
A. Linusanas     310.3047     M1831023     [13]     Algential, Jestuar       A. Linusanas     CZhao M405     M1110403     Presenti study     China, Yunnan       A. subrestas     CZLao 4221     M110403     Presenti study     China, Yunnan       A. subrestas     CLZao 4221     M1506313     Presenti study     China, Yunnan       A. subrestas     CLZao 4722     M5243499     Presenti study     China, Bejing       A. subrestas     CLZao 3832     M1114122     Urpublished     China, Yunnan       A. subrestas     CLZao 3852     M5243499     Presenti study     China, Bejing       A. subrestas     CLZao 3852     M5208444     Present study     China, Yunnan       A. subrestas     CLZao 3852     M5208444     Present study     China, Yunnan       A. correa     CDD 94255     KX221146     [23]     Thaliand, Chinag Reing       A. distat     CLZao 4035     M5243505     Present study     China, Yunnan       A. distat     CLZao 4030     M5243505     Present study     China, Yunnan       B. abasta     HHT-12226-Sp     K7134983     [25]     USA, Worcester       B. abasta     HHT-12226-Sp     K7134983     [25]     USA, Worcester       B. abasta     HHT-12226-Sp     K7134983     Present study <td>A. cerussatus</td> <td>CLZIII CLZIII CL</td> <td>MT2210E2</td> <td>[10]</td> <td>Arcontino Focuol</td>	A. cerussatus	CLZIII CL	MT2210E2	[10]	Arcontino Focuol		
A. Istatelinina     CLZID 34405     NExt-back     Predent Buddy     China, Belling       A. Istatelinina     CLZD 3421     Mik143168     Predent Buddy     China, Belling       A. mitrabilis     CLZDa 3422     Mik143168     Predent Study     China, Juman       A. subresus     ILZDa 3422     Mik34389     Predent Study     China, Vurnan       A. subresus     ILZDa 3422     Mik34389     Predent Study     China, Yurnan       A. subresus     ILEDA 3425     Mik104055     [21]     China, Yurnan       A. Intradiationa     CLZha 3562     Mik26844     Present Study     China, Yurnan       A. sisticia     OM 13932     Mik19318     [19]     China, Yurnan       A. corna     PDD 94825     KX621146     [23]     Thailand, Chinag Rai       A. corna     PDD 94825     KX621146     [23]     China, Yurnan       A. delicitat     MHU102112     KX621149     [23]     China, Yurnan       A. delicitat     MHU102112     KX621149     [23]     China, Yurnan       B. diato     OTina, Pinza     KX72129     [24]     China, Yurnan       B. diato     CLZha 4573     KY72279     [24]     China, Yurnan       B. diato     CLZha 4573     KY72279     [24]     China, Yurnan       B.	A. cerussulus	SFG 3004	NI1051050	[19] Dura ana tanta dari	China Vernan		
A. Issteptimics       The 224       PMT10903       [11]       China, Yunnan, Sequel         A. minibilis       MD2122       MC434489       Present study       China, Yunnan, Sequel         A. minibilis       MD222       MC434489       Present study       China, Yunnan         A. submers       He 4814       MH10905       Present study       China, Numan         A. submers       He 4814       MH10905       Present study       China, Yunnan         Attricidurin sistica       CLZhao 3652       ME268844       Present study       China, Numan         A. contei       CLZhao 4245       MK243930       Present study       China, Yunnan         A. contei       CLZhao 4245       MK24395       Present study       China, Yunnan         A. dicitati       CLZhao 4245       KK21146       [23]       Thaliand, Ching, Rai         A. dicitati       CLZhao 4310       MK34395       Present study       China, Yunnan         B. dicitati       HH119226-Sp       KF134983       [25]       USA, Worcester         B. dicitati       HH212226-Sp       KF134983       [25]       USA, Worcester         B. dicitati       HH212226-Sp       KF134980       [27]       USA, Worcester         Cantabarolas cibritis       CL4	A. isabellinus	CLZna0 4405	NIK343482	Present study	China, Yunnan		
A. minimis         CL2.00 4021         M2542488         Present study         China, Iuman           A. subress         CL2hao 4722         MK34849         Present study         China, Yunnan           A. subress         CL2hao 4722         MK34849         Present study         China, Yunnan           A. subress         CL2hao 4722         MK34849         Present study         China, Yunnan           A. subress         CL2hao 4562         MK29844         Present study         China, Yunnan           A. arcial static         CL2hao 4562         MK20844         Present study         China, Yunnan           A. arcial static         CL2hao 4245         MK20844         Present study         China, Yunnan           A. arcial static         CL2hao 4245         KK24844         Present study         China, Yunnan           A. arcial static         CL2hao 4310         KK343050         Present study         China, Yunnan           A. adictat         CL2hao 4310         MK343058         Present study         China, Yunnan           B. adictat         CL2hao 4310         MK343059         Present study         China, Yunnan           B. adictat         CL2hao 4318         MT17289         Present study         China, Yunnan           B. adictat         CL2hao 4303<	A. isubellinus	He 3294	MH109055	[21] Due a sur tratta das	China, beijing		
A. miranusJub 2152M1 250513[19]Argentina, FsqluetA. subreseusHe 4814MF1100055[21]China, NunnanAutroida malicolaCLZhao 3822MH114612UpublishedChina, YunnanAutroida residueSFC 20180314-01MK592816[22]Korea, ScoulA. corneaCLZhao 3552MK208831[19]China, BeijingA. corneaCLZhao 4245MK043502Present studyChina, YunnanA. corneaCLZhao 4245KK021145[23]Thailand, Chinag BaiA. corneaHDD 94625KK021145[23]Thailand, Chinag BaiA. corneaHDD 94625KK021145[23]Thailand, Chinag BaiA. defactaMKU1626373KK72229[23]Thailand, Chinag BaiA. defactaMKU1626373KK72229[23]Thailand, Chinag BaiB. defactaCLZhao 9318MT177289Present studyChina, YunnanB. defactaCLZhao 9318MT177289Present studyChina, YunnanB. defactaCLZhao 9318MT177289Present studyChina, YunnanC. corninCLZhao 9318MT177289Present studyChina, YunnanC. corninCLZhao 9318MT177289Present studyChina, YunnanC. corninCLZhao 9302KC88532Present studyChina, YunnanC. corninCLZhao 9379MK20856Present studyChina, YunnanC. corninCLZhao 9379MK208582Present studyChina, YunnanC. cornin	A. mirabilis	CLZhao 4621	MIK343488	Present study	China, Yunnan		
A. statisticsCL2.Aux 92/2IAK 34389Present studyChina, TunnanA. statisticsC. Likuo 383MH110615[21]China, TunnanAntrodu malicoluCL2.Huo 3832MH114612UupublishedChina, YunnanAntrodu malicoluCL2.Huo 3852MK208881Present studyChina, YunnanA. acaticaCL2.Huo 3552MK208881Present studyChina, YunnanA. acaticaCL2.Huo 1245MK208881Present studyChina, YunnanA. acaticaCL2.Huo 1245KK621146[23]China, YunnanA. acaticaCL2.Huo 1403KK201146[24]China, YunnanA. deficitaMT11016112KK201149[23]China, YunnanA. deficitaCL2.huo 4310MK343505Present studyChina, YunnanB. adistaHHB-1280-59RV134983[25]USA, WorcesterBodetus adultis375FU556644[26]Spain, CabrilsBesomentalitis cortuinCL2.huo 4308MK228856Present studyChina, YunnanB. cortuinFP-102382KV135007[25]USA, WorcesterCamboritis cortuinCL2.huo 3603MK288869Present studyChina, YunnanC. acataCL2.huo 4931MK288860Present studyChina, YunnanC. acataCL2.huo 4937MK288860Present studyChina, YunnanC. acataCL2.huo 4937MK268869Present studyChina, YunnanC. acataCL2.huo 4937MK268860Present studyChina, Yun	A. mirabilis	JMB 2152	M1568513		China Vumpan		
A. stutbestesHe 4814MH109055[21]Chrina, BeijingAttrodit asiditicaCIZhao 3822MH114612UpublishedChina, YunnanAttrodit asiditicaSFC 20180314-01MK992816[22]Koree, SeoulA. asiditicaOM 13932MZ618931[19]China, BeijingA. corneaCIZhao 3652MK26884Present studyChina, WinnanA. corneaFDD 94825KX261146[23]Thailand, Chiang RaiA. corneaFDD 94825KX261146[23]Thailand, Chiang RaiA. delicatuMFU106112KX261149[23]Thailand, Chiang RaiA. adictatMFU106112KX261149[23]Thailand, Chiang RaiA. adictatMFU106112KX261149[23]UbA, WorcesterB. datastaCIZhao 6310MK343509Present studyChina, YunnanB. datastaCIZhao 6310MK743509Present studyChina, YunnanB. datastaFHB 12826-5pEH139686Present studyChina, YunnanB. datastaFHB 2326-5pKH135007[26]Candrolis AbrinisB. datastaFHB 2326-5pKH23804[26]Spain, CabrilaB. datastaFHB 2326-5pKH23804[26]Candrolis AbrinisCandrolis abrinisFD 2653217MK298866Present studyChina, YunnanCandrolis abrinisCLAo 3093MK298866Present studyChina, YunnanCandrolis abrinisCLAo 3502MK298856Present studyChina, YunnanC	A. subroseus	CLZhao 4722	MK343489	Present study	China, runnan		
Arthrodu maticulaCL2hao 3832MH114612UnpublishedChina, YunnanArticularia asiaticaCL2hao 3652MK268844Present studyChina, SeuilaA. corieaCLZhao 4245MK34502Present studyChina, YunnanA. corieaCLZhao 4245MK34502Present studyChina, WannanA. corieaCLZhao 4245MK34505Present studyChina, WannanA. corieaCLZhao 4403MK343505Present studyChina, BeijingA. delicataCLZhao 4403MK343506Present studyChina, BeijingBirkmdiera adustaCLZhao 4310MK343508Present studyChina, YunnanB. datistaHHE-1282c-SpKP134983[25]USA, WorcesterBolctas dalisis375FU554664[26]Spain, CabrilsB. coriumCLZhao 9318MT17289Present studyChina, YunnanB. coriumCL2hao 9318MT12807[27]Canada, Prince GeorgeCartipari lacerataCL2hao 9318MT268865Present studyChina, YunnanC. lacerataFP55521TKP33024[25]USA, WorcesterCartibarellis cibririsCL-67MK26888Present studyChina, YunnanC. lacerataFP55521TKP35024[23]USA, WorcesterCartistar moscarmarmeDai 12892KK26887Present studyChina, YunnanC. lacerataCL2hao 3370MK268888Present studyChina, YunnanC. labecimamoneaDai 12892KK26857Present study<	A. subroseus	He 4814	MH109055	[21]	China, Beijing		
Abeta epiptyliaSEC 2108(314-01MK 992816[22]Korea, ScoulA. arisiticaOM 13932MK268844Present studyChina, BeijingA. coristaCLZbao 4245MK343502Present studyChina, BeijingA. coristaPDD 94825KK2621146[23]Thailand, Chiang BaiA. delicitaMFU1162112KK2621146[23]Thailand, Chiang BaiA. delicitaMFU1162112KK2621149[23]Thailand, Chiang BaiA. delicitaMFU1162112KK2621149[23]Thailand, Chiang BaiB. abustaCLZbao 4310MK343508Present studyChina, YunnanB. abustaCLZbao 4310MK343508Present studyChina, YunnanB. abustaCLZbao 9318MT172799Present studyChina, YunnanB. coriumFP-102382KP134983[23]U5A, WorcesterBueletus edulis375FU53664[23]U5A, WorcesterCamburellus coinrineCLZbao 3693MK268856Present studyChina, YunnanB. coriumFP-102382KP135007[23]U5A, WorcesterCamburellus coinrineCLZbao 3693MK268856Present studyChina, YunnanC. iaccerataCLZbao 3602MK268858Present studyChina, YunnanC. iaccerataCLZbao 3502MK268858Present studyChina, YunnanC. corataCLZbao 4379MK268858Present studyChina, YunnanC. corataCLZbao 4479MK268858Present studyChina, Yu	Antrodia malicola	CLZhao 3832	MH114612	Unpublished	China, Yunnan		
Antricularia assistica     CL2ao 3652     MZ68984     Present study     China, Wunnan       A. cornea     CLZao 4245     MK343502     Present study     China, Yunnan       A. cornea     PDD 94825     KX621146     [23]     Thailand, Chinag Rai       A. delicata     CLZao 4403     MK343505     Present study     China, Yunnan       A. delicata     CLZao 4403     MK343505     Present study     China, Yunnan       A. delicata     CLZao 4403     MK343505     Present study     China, Yunnan       B. daista     HH=12826-Sp     KYA21148     [23]     USA, Worcester       Boltus dulits     375     EU354664     [26]     Spain, Cabrils       B. adusta     HH=12826-Sp     KP133007     [23]     USA, Worcester       Curtural is colurius     CL2hao 3918     MT17289     Present study     China, Yunnan       B. acrium     CL2hao 3693     MK264856     Present study     China, Yunnan       C. accrata     FP-555217     KP38024     [25]     USA, Worcester       Carloria iaccrata     CL2hao 3802     MK264858     Present study     China, Yunnan       C. accrata     FP-555217     KP38024     [25]     USA, Worcester       C. accrata     FP-555217     KP38024     [26]     USA, Worcester <td>Athelia epiphylla</td> <td>SFC 20180314-01</td> <td>MK992816</td> <td>[22]</td> <td colspan="2">Korea, Seoul</td>	Athelia epiphylla	SFC 20180314-01	MK992816	[22]	Korea, Seoul		
A. astaticaOM 13932MI2518931[19]China, BeijingA. corneaPDD 94825KX621146[23]Thailand, Chiang RaiA. delicataClZhao 4403MK543505Present studyChina, DeijingA. delicataMFU162112KX621149[23]Thailand, Chiang RaiA. delicataMFU162112KX621149[23]Thailand, Chiang RaiA. mesentricaBRNM 64873KP22279[24]China, DeijingBjerkmidera adustaCLZhao 4310MK543508Present studyChina, VinnanB. adustaHIB 12865 PKP134983[25]USA, WorcesterBoletus edulis375EU554664[26]Spain, CabrilsByssemeritisc coriumFP102382KP135007[25]USA, WorcesterConlinentilis coriumFP102382KP135007[25]USA, WorcesterCentharellus cibariusCL-67MK268856Present studyChina, YunnanC. lacerataCLZhao 3693MK268856Present studyChina, VunnanC. alocinnamoneaCLZhao 4193MK268860Present studyChina, DeijingC. c. alocinnamoneaCLZhao 4193MK268860Present studyChina, VunnanC. c. alocinnamoneaCLZhao 4300MK268866[19]Netherlands, UpsalalaanC. c. alocinnamoneaCLZhao 4320MK268877Present studyChina, YunnanC. c. andaCLZhao 4320MK268870[19]Netherlands, UpsalalaanC. c. dropserzesKMRB 18041009MK120528Pr	Auricularia asiatica	CLZhao 3652	MK268844	Present study	China, Yunnan		
A. corneaCL2hao 2425MK343502Present studyChina, YunnanA. delicataCL2hao 4403MK343505Present studyChina, YunnanA. delicataCL2hao 4403MK343505Present studyChina, YunnanA. delicataMFU162112KX621149[23]Thailand, Chiang RaiA. mesentericaBRNM 648573KP229279[24]China, YunnanB. adustaHHD-12826-SpKP134983[25]USA, WorcesterBoletus edulis375EU354664[26]Spain, CabrilsB. coriumCL2hao 9318MT177289Present studyChina, YunnanB. coriumFP-102882KP135007[25]USA, WorcesterCambardius cibrariusCL-67MK281467[27]Canada, Prince GeorgeCaribroi lacerataCL2hao 3693MK268860Present studyChina, YunnanC. aluerataCL2hao 3502MK268860Present studyChina, YunnanC. aluerataCL2hao 3502KK485522[19]China, YunnanC. andatIL2Mao 3502MK268860Present studyChina, YunnanC. andataCL2hao 3502MK268863[20]Netherlands, UpsalalaanC. contataCL2hao 3502MK268867Present studyChina, YunnanC. contataIL2Mao 3502MK268877Present studyChina, YunnanC. corestmCL2hao 3504MK20528[19]Netherlands, UpsalalaanC. rosteumCLS140MK243523Present studyChina, YunnanC.	A. asiatica	OM 13932	MZ618931	[19]	China, Beijing		
A. cornuePDD 94825KX621146[23]Thailand, Chiang RaiA. delicataMFLU162112KX621149[23]Thailand, Chiang RaiA. desentericaBRNM 64873KY2229[24]China, WunnanB. A. dustaCLZhao 4310MK343508Present studyChina, DeijingBjerkandera adustaCLZhao 4310MK343508Present studyChina, WunnanB. adustaHIB-12826SpKP134983[25]USA, WorcesterBoletus edulis375FU1554644[26]Spain, CabrilsB. cariumF2-10282KP135007[25]USA, WorcesterCantharellus cibriusCLTao 3918MT177289Present studyChina, YunnanC. claratiaF12-10282KP135007[25]USA, WorcesterCarinotia lacentaCLZhao 693MK268856Present studyChina, YunnanC. lacentiaF12-5521TKP135024[25]USA, WorcesterCremen alborinamomeaDai 12892KC485522[19]China, YunnanC. albocinnamomeaDai 12892KC485563Present studyChina, YunnanC. cubcinnamomeaDai 12892KC485564[20]Netherlands, UpsalalaanC. cubcinnamomeaDai 12592MK268860[19]Netwarade, UpsalalaanC. cubcinnamomeaCLZhao 3562MK268563[20]Netherlands, UpsalalaanC. cubcinnamomeaCLZhao 4379MK243524Present studyChina, YunnanC. cubcinnamomeaCLZhao 4379MK243524Present study <td< td=""><td>A. cornea</td><td>CLZhao 4245</td><td>MK343502</td><td>Present study</td><td>China, Yunnan</td></td<>	A. cornea	CLZhao 4245	MK343502	Present study	China, Yunnan		
A. delicataCLZhao 4403MK243505Present studyChina, XunnanA. mesentericaBRNM 648573KP220279[24]China, BelingA. mesentericaBRNM 648573KP220279[24]China, VunnanBjerkander adustaHHB-12826-SpKP134983[25]USA, WorcesterBoletus edulis375EU554664[25]Spain, CabrilsBysomerulius coriumC1Zhao 9318MT17289Present studyChina, YunnanB. coriumFP-10282KP135007[25]USA, WorcesterCentimelius cibariusC1Zhao 3693MK268856Present studyChina, YunnanC. lacerataC1Zhao 3693MK268856Present studyChina, WinnanC. lacerataC1Zhao 3693MK268860Present studyChina, BeljingC. albocinnamoneaC1Zhao 4193MK268860Present studyChina, WinnanC. albocinnamoneaC1Zhao 3502MK268852[19]China, WinnanC. c.abocinnamoneaC1Zhao 4193MK268860Present studyChina, WinnanC. c.abocinnamoneaC1Zhao 4193MK268873Present studyChina, WinnanC. c.abocinnamoneaC1Zhao 4193MK268874[20]Netherlands, UpsalalaanC. constaC1Zhao 4137MK268877Present studyChina, YunnanC. constaC1Zhao 4379MK345324Present studyChina, YunnanC. raveournumCBS 330.66MH858610[20]Netherlands, UpsalalaanC. raveournumCBS 330.66MH858674	A. cornea	PDD 94825	KX621146	[23]	Thailand, Chiang Rai		
A. delicataMFLU162112KX621149[23]Thailand, Chiang RaiA. mesutericaBRNM 64873KY2279[24]China, BejingBiprknudera adustaCLZhao 4310MK343508Present studyChina, YunnanB. adustaHTB-12826-5pKP1349433[25]USA, WorcesterBoletus eduits375FU534664[26]Spain, CabrilsBysomerrilis cortumCLZhao 318MT177289Present studyChina, YunnanB. cariumFP-102382KP135007[27]Canda, Prince GeorgeCartharellus cibariusCL-67MK281467[27]Canda, Prince GeorgeCariperia lacerataFP-55521TKP135024[23]USA, WorcesterC. recena discrimanomaO tal 12892KC48855Present studyChina, YunnanC. albocinnanomaO tal 12892KC48852[19]China, MinanaC. zonutaICLPha 3502MK268860Present studyChina, YunnanC. albocinnanomaO tal 2892KC488572Present studyChina, YunnanC. albocinnanomaCLZhao 4379MK343524Present studyChina, YunnanC. rosecarneumCLZhao 4379MK343524Present studyChina, YunnanC. rosecarneumCLZhao 4379MK2485074[20]Netherlands, UpsalalanC. rosecarneumCLZhao 4420MK26786[19]Netherlands, UpsalalanC. rosecarneumCLZhao 4420MK26786[20]Netherlands, UpsalalanC. rosecarneumCLZhao 4426MK26790 <td>A. delicata</td> <td>CLZhao 4403</td> <td>MK343505</td> <td>Present study</td> <td>China, Yunnan</td>	A. delicata	CLZhao 4403	MK343505	Present study	China, Yunnan		
A. mesentericaBRNM 648573KP29279[24]China, BeijingBirkmder adustaHHB-12826-SpKP134983[25]USA, WorcesterBoltus edulis375EUS54664[26]Spain, CabrilsBysomerulius coriumCLZhao 9318MT177289Present studyChina, YunnanB. coriumFP-102382KP135007[25]USA, WorcesterCantharefilius cibariusCL-67MK281467[27]Canada, Prince GeorgeCeriporia lacerataCLZhao 3693MK268856Present studyChina, YunnanC. lacerataFP-55521TKP135024[25]USA, WorcesterC. allocinnamomeaCLZhao 3502MK268860Present studyChina, YunnanC. allocinnamomeaCLZhao 3502KK268522[19]NetwesterC. cutataICM-6437MW2688680Present studyChina, BujingC. cutataICM-6437MW2688638Present studyChina, VunnanC. cutataICM-6437MW856063[20]Netherlands, UpsalalaanC. paideherimusCBS 130.40MH856063[20]Netherlands, UpsalalaanC. rascarneumCLZhao 3502MH268877Present studyChina, YunnanC. rascarneumCLZhao 3562MH26874[23]Netherlands, UpsalalaanC. rascarneumCBS 330.66MH85810[20]Netherlands, UpsalalaanC. rascarneumCLZhao 3562MH26790[24]USA, BethesdaD. sterologisCutaba 433MK269228Present studyChina,	A. delicata	MFLU162112	KX621149	[23]	Thailand, Chiang Rai		
Bjerkandera adustaCLZhao 4310MK343508Present studyChina, YunnanB. adustaHHB-12826-5cpKP134983[25]USA, WorcesterBoletus eduitis375EU554664[25]Spain, CabrilsBysomerulius coriumCLZhao 9318MT177289Present studyChina, YunnanB. coriumFP-102382KP135007[25]USA, WorcesterCanthareflus cibariusCL-67MK281467[27]Canada, Prince GeorgeCritioni lacerataFP-55521TKP135024[25]USA, WorcesterCerreau albocinnamomeaCLZhao 4193MK268860Present studyChina, YunnanC. albocinnamomeaDati 12892KC485522[19]China, YunnanC. albocinnamomeaDati 12892KC485523[19]New Zealand, AucklandC. autatICMP 16347MW862786[19]New Zealand, AucklandC. anataICMP 16347MW862786[19]Netw Zealand, AucklandC. furicium roseocarneumCBS 30.66MH858810[20]Netherlands, UpsalalaanC. roseiumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseiumCBS 104.52MH856944[20]	A. mesenterica	BRNM 648573	KP729279	[24]	China, Beijing		
B adutsaHHB-12826-SpKP134983[25]USA, WorcesterBoletus adutis375FUIS54664[26]Spain, CabrilsByssomerulius coriumCLZnao 9318MT177289Present studyChina, YunnanB. coriumFP102382KP135007[25]USA, WorcesterCantharellus cibariusCL-67MK281467[27]Canada, Prince GeorgeCariporia lacerataCLZnao 3693MK268886Present studyChina, YunnanC. laccrataFP55521TKP13024[25]USA, WorcesterC. allocinnamomeaCLZnao 4193MK268860Present studyChina, BejingC. c. allocinnamomeaDai 12892KC485522[19]China, BejingC. zonataCLZnao 3502MK268838Present studyChina, YunnanC. zonataCLZnao 4379MK343524Present studyChina, YunnanC. cnicherrimusCDS 130.40MH856063[20]Netherlands, UpsalalaanC. ruscomneumCDS 330.66MH458810[20]Netherlands, UpsalalaanC. ruscomneumCDS 104.52MH250944[20]Netherlands, UpsalalaanC. ruscomneumCDS 104.52MH250944[20]Netherlands, UpsalalaanC. ruscomneumCDS 104.52MH250944[20]Netherlands, UpsalalaanC. ruscomneumCDS 104.52MH250944[20]Netherlands, UpsalalaanC. ruscomneumCDS 104.52MH26928Present studyChina, YunnanC. ruscomneumCDS 104.52MH269944[20]<	Bjerkandera adusta	CLZhao 4310	MK343508	Present study	China, Yunnan		
Boletus adults375FUS54664[26]Spain, CabrilsBysomerulius coriumCLZao 9318MT177289Present studyChina, YunnanB. coriumFP-102382KP135007[25]USA, WorcesterCamlanellus cibariusCL-67MK281467[27]Canada, Prince GeorgeCerripoit lacerataCLZhao 3693MK268856Present studyChina, YunnanC. lacerataFP-55521TKP135024[25]USA, WorcesterCerreia alcoriumamoneaCLZhao 4193MK268860Present studyChina, YunnanC. albocinnamoneaCLZhao 3502MK268863[19]New Zealand, AucklandC. conataICMP 16347MW862786[19]New Zealand, AucklandClimacodon pulcherrimusCDS 130.40MH856063[20]Netherlands, UpsalalaanC. pulcherrimusCLZhao 4379MK343524Present studyChina, YunnanC. croscumCB 30.66MH858810[20]Netherlands, UpsalalaanC. roscumCB 104.52MH850644[20]Netherlands, UpsalalaanC. roscumCB 104.52MH850647[29]Korea, SeoulDaraymeye syntexyCLZhao 4143MK262288Present studyChina, YunnanC. chrysocreasKMRB 18041009MK120528[19]Korea, SeoulD. stillatusUPS F-939814MN595677[30]Sweden, UpsalaD. datelliformisPDD 76666NR.166790[29]USA, BethesdaD. stillatusUPS F-939814MK33522Present study	B. adusta	HHB-12826-Sp	KP134983	[25]	USA, Worcester		
Byseomenulius coriumCLZhao 9318MT177289Present studyChina, YunnanB. coriumFP-10282KP135007[25]USA, WorcesterCanthanellus cibariusCL-67MK281467[27]Canada, Prince GeorgeCeriporia lacentaCLZhao 3693MK268856Present studyChina, YunnanC. lacentaFP-55511KP135024[25]USA, WorcesterCentronamomeaDai 12892KC485522[19]China, BeijingC. contaCLZhao 3702MK268838Present studyChina, YunnanC. contaCLZhao 3502MK268838Present studyChina, YunnanC. contaCLZhao 3502MK268838Present studyChina, YunnanC. contaCLZhao 3502MK268837Present studyChina, YunnanC. rosecurneumCBS 130.40MH856063[20]Netherlands, UpsalalaanC. rosecurneumCLZhao 3562MK268877Present studyChina, YunnanC. rosecurneumCLZhao 3562MK26879Present studyChina, YunnanC. chrysocrasCLZhao 4143MK20528Present studyChina, YunnanC. chrysocrasCLZhao 4143MK20528[19]USA, BethesdaD. fidelliformisPDD 107980NR, 146790[23]USA, BethesdaD. fidelliformisPDD 76696NR, 166790[29]USA, BethesdaD. fidelliformisCLZhao 4260MK343522Present studyChina, YunnanD. sterooidesCLZhao 4377MT93539Present study <t< td=""><td>Boletus edulis</td><td>375</td><td>EU554664</td><td>[26]</td><td>Spain, Cabrils</td></t<>	Boletus edulis	375	EU554664	[26]	Spain, Cabrils		
B. cortiumFP-102382KP135007[25]USA, WorcesterCantharellus cibariusCL-67MK281467[27]Canada, Prince GeorgeCarriporia lacentaCLZhao 3693MK268886Present studyChina, YunnanC. lacerataFP-55521TKP135024[25]USA, WorcesterCerrena albocinnuamoneaCLZhao 4193MK268886Present studyChina, YunnanC. albocinnuamoneaCLZhao 3502KC485522[19]China, BeijingC. zonataICMP 16347MW862786[19]New Zealand, AucklandC. pulcherrinnusCLZhao 4379MK343524Present studyChina, YunnanC. conataICMP 16347MW862786[20]Netherlands, UpsalalaanC. conataICMP 16347MW862786[20]Netherlands, UpsalalaanC. concarneumCLZhao 4379MK343524Present studyChina, YunnanC. concarneumCLZhao 3562MK268877Present studyChina, YunnanC. coscurneumCLZhao 3562MH856944[20]Netherlands, UpsalalaanC. croscurneumCLZhao 4143MK26928Present studyChina, YunnanC. chirgocreasCLZhao 4143MK269577[30]Sweden, UpsalaD. fidaleliformisPDD 107980NR,148190[28]USA, BethesdaD. diadlopsis confragosaCLZhao 4260MK34522Present studyChina, YunnanD. confragosaCLZhao 4260MK345323Present studyChina, YunnanD. sereoidesCLZhao 4372 <td>Byssomerulius corium</td> <td>CLZhao 9318</td> <td>MT177289</td> <td>Present study</td> <td>China, Yunnan</td>	Byssomerulius corium	CLZhao 9318	MT177289	Present study	China, Yunnan		
Gantharellus cibriusCL-67MK281467[27]Canada, Prince GeorgeCeriporia lacentaCL3ao 3693MK268856Present studyChina, YunnanC. lacentaFP-555211KP135024[25]USA, WorcesterCertena albocinnamoneaDai 12892KC485522[19]China, YunnanC. albocinnamoneaDai 12892KC485522[19]China, BeijingC. contatCLZhao 3502MK268838Present studyChina, BeijingC. contatICMP 16347MK862786[19]New Zealand, AucklandC. pulcherrimusCLB 10.00MH856063[20]Netherlands, UpsalalaanC. pulcherrimusCLZhao 4379MK345324Present studyChina, YunnanC. rosecumCBS 330.66MH856044[20]Netherlands, UpsalalaanC. rosecumCLZhao 3562MK268877Present studyChina, YunnanC. rosecumCLZhao 514.52MH856944[20]Netherlands, UpsalalaanC. crosecumCLZhao 1413MK205228Present studyChina, YunnanC. crosecumCLZhao 1413MK205228[19]Korea, SeoulDacrymyces cyrtosporusPDD 107980NR, 148190[25]USA, BethesdaD. fidelliformisPD 76696NR, 166790[29]USA, BethesdaD. fidelliformisPD 76696NR, 166790[29]China, YunnanD. corifragosaCLZhao 4220MK343523Present studyChina, YunnanD. drifragosaCLZhao 4377MT93539Present stud	B. corium	FP-102382	KP135007	[25]	USA, Worcester		
Ceriporia lacerataCL2hao 3693MK268856Present studyChina, YunnanC. lacerataFP-55521TKP135024[25]USA, WorcesterCerrena albocinnamomeaCL2hao 4193MK268860Present studyChina, YunnanC. albocinnamomeaDai 12892KC485522[19]China, YunnanC. zonataCL2hao 3502MK26883Present studyChina, YunnanC. zonataICMP 16347MW862786[19]New Zealand, AucklandC. pulcherrimusCL2hao 4379MK343524Present studyChina, YunnanC. pulcherrimusCL2hao 3502MK843524Present studyChina, YunnanC. pulcherrimusCL2hao 4379MK343524Present studyChina, YunnanC. rosecarneumCL2hao 3562MK848587Present studyChina, YunnanC. rosecarneumCBS 104.52MH856944[20]Netherlands, UpsalabaanC. crosecarneumCBS 104.52MH856944[20]Netherlands, UpsalabaanC. crosecarneumCBS 104.52[19]Korea, ScoulD. fabelliformisPDD 107980NR, 166790[29]USA, BethesdaD. fabelliformisPDD 76696NR, 166790[29]USA, BethesdaD. stillatusUPS F-939814MK543522Present studyChina, YunnanD. confragosaCu1283KU900526[19]China, YunnanD. confragosaCu1283KU900526[19]China, YunnanD. steroidesCU2hao 4320MK34352Present studyChina,	Cantharellus cibarius	CL-67	MK281467	[27]	Canada, Prince George		
CLacerataFP-55521TKP135024[25]USA, VorcesterCernera albocinnamomeaDai 12892KC485522[19]China, VunnanC. albocinnamomeaDai 12892KC485522[19]China, BeijingC. zonataCLZhao 3502MK268838Present studyChina, YunnanC. zonataICMP 16347MW862786[19]New Zealand, AucklandClimacodon pulcherrimusCBS 130.40MH856063[20]Netherlands, UpsalalaanC. pulcherrimusCLZhao 4379MK343524Present studyChina, YunnanC. rosecurneumCBS 30.66MH858810[20]Netherlands, UpsalalaanC. rosecurneumCBS 30.66MK268877Present studyChina, YunnanC. rosecurneumCBS 104.52MK268877Present studyChina, YunnanC. rosecurneumCBS 104.52MK26928Present studyChina, YunnanC. rosecuresCLZhao 34143MK269228Present studyChina, YunnanC. chrysocrasCLZhao 4143MK269228[19]Korea, SeoulDacrymcycs cyrtosporusPDD 107980NR 148190[28]USA, BethesdaD. flabeliformisPDD 76696NR 164790[29]USA, BethesdaD. steroidesCui 2883KU900526[19]China, YunnanD. confragosaCui 2843KU900526[19]China, YunnanD. steroidesCui 8132JX59270[19]China, YunnanD. steroidesCui 8132JX59920[19]China, Yunnan <td>Ceriporia lacerata</td> <td>CLZhao 3693</td> <td>MK268856</td> <td>Present study</td> <td>China, Yunnan</td>	Ceriporia lacerata	CLZhao 3693	MK268856	Present study	China, Yunnan		
Cerrena albocinnamomeaCLZhao 1193MK268860Present studyChina, YunnanC. albocirnamomeaDai 12892KC485522[19]China, WinnanC. zonataCLZhao 3502MK268838Present studyChina, WinnanC. zonataICMP 16347MW862786[19]New Zealand, AucklandClimacodon pulcherrimusCLZhao 4379MK343524Present studyChina, YunnanC. rosecarneumCLZhao 350.66MH858810[20]Netherlands, UpsalalaanC. rosecarneumCLZhao 3562MK268877Present studyChina, YunnanC. rosecarneumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. rosecarneumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseumCBS 104.52MH856974[20]Netherlands, UpsalalaanC. drigsocrasCLZhao 4143MK26228Present studyChina, YunnanC. drigsocrasPDD 107980NR, 148190[28]USA, BethesdaD. fabelliformisPDD 76696NR, 166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDacrymyces cyrlososaCui 2883KU900526[19]China, YunnanD. stroidesCLZhao 9427MK543539Present studyChina, YunnanD. stroidesCLZhao 5722MK268896[19]China, YunnanD. stroidesCui 2833JK559270 <td< td=""><td>C. lacerata</td><td>FP-55521T</td><td>KP135024</td><td>[25]</td><td>USA, Worcester</td></td<>	C. lacerata	FP-55521T	KP135024	[25]	USA, Worcester		
C. albocimumomeaDai 12892KC485522[19]China, BeijingC. zonataCLZhao 3302MK268838Present studyChina, YunnanC. zonataICMP 16347MW862786[19]New Zealand, AucklandClinacodon pulcherrimusCBS 130.40MH856063[20]Netherlands, UpsalalaanC. pulcherrimusCLZhao 4379MK343524Present studyChina, YunnanCarticium roseocarneumCBS 330.66MH858810[20]Netherlands, UpsalalaanC. rosecumCLZhao 3562MK268877Present studyChina, YunnanC. rosecumCLZhao 3562MK269228Present studyChina, YunnanC. crosecumCLZhao 4143MK269228[19]Korea, SeoulDarrymyces cyrtosporusPDD 107980NR,148190[28]USA, BethesdaD. flabelljornnisPDD 76666NR,166790[29]USA, BethesdaD. flabelljornnisPDD 76666NR,166790[29]USA, BethesdaD. orifragosaCLZhao 4260MK343523Present studyChina, YunnanD. confragosaCLZhao 4260MK343523Present studyChina, YunnanD. stereoidesCLZhao 9377MT193539Present studyChina, YunnanD. stereoidesCLZhao 9372KK268856[19]China, YunnanD. stereoidesCLZhao 3722KK268856[19]China, YunnanE. scabrosaCLZhao 3722KK268856[19]China, YunnanF. anbrosiusCLZhao 3722KK268896Present stu	Cerrena albocinnamomea	CLZhao 4193	MK268860	Present study	China, Yunnan		
C. zonataC. LZhao 3502MK268838Present studyChina, YumanC. zonataICMP 16347MW862786[19]New Zealand, AucklandClinacodon pulcherrinnusCLZhao 4379MK343524Present studyChina, YunnanC. pulcherrinnusCLZhao 4379MK343524Present studyChina, YunnanCorticium roseocarneumCLZhao 3562MK248877Present studyChina, YunnanC. rosecarneumCLS 330.66MH858810[20]Netherlands, UpsalalaanC. rosecarneumCLS 430.65MK26828Present studyChina, YunnanC. rosecumCB 104.52MH856944[20]Netherlands, UpsalalaanC. crosecarneumCLS 44143MK26928Present studyChina, YunnanC. chrysocrasKMRB 18041009MK120528[19]Korea, SeoulDarymyccs cyrtosporusPDD 176696NR, 148190[28]USA, BethesdaD. fidbellformisPDD 107980NR, 148190[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDatornia mollisCLZhao 4260MK343523Present studyChina, YunnanD. sonfragosaCui 283KU900526[19]China, BeijingDatronia mollisCLZhao 4259MK343523Present studyChina, YunnanD. steroidesCLZhao 937MT193539Present studyChina, YunnanD. steroidesCLZhao 3722MK268896Present studyChina, BeijingFarileila scabrosaCui 6236	C. albocinnamomea	Dai 12892	KC485522	[19]	China, Beijing		
C. zonataICMP 16347MW862786[19]New Zealand, AucklandClimacodon pulcherrinusCBS 130.40MH856063[20]Netherlands, UpsalalaanC. pulcherrinusCLZhao 4379MK343524Present studyChina, YunnanCorticium roseocarneumCBS 330.66MH858810[20]Netherlands, UpsalalaanC. rosecoarneumCLZhao 3562MK268877Present studyChina, YunnanC. rosecoarneumCLZhao 4143MK269228Present studyChina, YunnanC. crosetumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. rosecoarneuerPDD 107980NR 148190[28]USA, BethesdaD. flabelliformisPDD 107980NR 148190[28]USA, BethesdaD. flabelliformisPDD 76696NR 166790[29]USA, BethesdaD. stillatusUPS F-39814MN595677[30]Sweeden, UppsalaDacargosaCu12Aa 4260MK343522Present studyChina, YunnanD. confragosaCu2Lhao 4220MK343523Present studyChina, PuijngDatronia mollisCLZhao 4220MK34553[19]China, GeoulD. stereoidesCLZhao 4220MK345254[19]Korea, SeoulD. stereoidesCLZhao 3722KJ668554[19]China, YunnanE acabrosaCLZhao 3722MK34896Present studyChina, YunnanE scabrosaCLZhao 3341MH114652Present studyChina, YunnanF anbosiusHulcr 12546LC216231[32	C. zonata	CLZhao 3502	MK268838	Present study	China, Yunnan		
Climacodon pulcherrimusCBS 130.40MH850063[20]Netherlands, UpsalalaanC. pulcherrimusCLZhao 4379MK343524Present studyChina, YunnanCorticium roseocarneumCLS 33.066MH858810[20]Netherlands, UpsalalaanC. roseocarneumCLZhao 3562MK268877Present studyChina, YunnanC. roseoumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseoumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseoumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. croseoumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. croseoumCBS 104.52MH856977Present studyChina, YunnanC. chrysocreasKMRB 18041009MK120528[19]Korea, SeoulD. fabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDadaleopsis confragosaCui 2883KU900526[19]China, YunnanD. otifragosaCui 2883KU900526[19]China, YunnanD. stereoidesCLZhao 4220MK343523Present studyChina, YunnanD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCLZhao 3722MK268966[19]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. sabrosaCui 2393MK343528	C. zonata	ICMP 16347	MW862786	[19]	New Zealand, Auckland		
C. pulcherrimusC.LZhao 4379MK343524Present studyChina, YunnanCorticium roseccarneumCLZhao 3562MK268877Present studyChina, YunnanC. roseumCBS 104.52MH856914[20]Netherlands, UpsalalaanC. roseumCLZhao 3562MK268877Present studyChina, YunnanC. roseumCLZhao 4143MK269228Present studyChina, YunnanC. chrysocreasKLMB 18041009MK120528[19]Korea, SeoulDacrymyces cyrtosporusPDD 107980NR_148190[28]USA, BethesdaD. fiabelliformisPDD 7666NR_166790[29]USA, BethesdaD. fiabelliformisPDD 7666NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCuZhao 4260MK343523Present studyChina, YunnanD. steroidesCLZhao 9937MT193539Present studyChina, YunnanD. steroidesCui 8132JX559270[19]China, BeijingD. steroidesCui 823KS6896Present studyChina, YunnanE. scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCLZhao 3721MK268896Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3411MH114652Pre	Climacodon nulcherrimus	CBS 130.40	MH856063	[20]	Netherlands, Upsalalaan		
ContinuityCLB 30.66MH258210FlortnethyNetherlands, UpsalalaanCorticium roseocarreumCLB 30.66MH258810[20]Netherlands, UpsalalaanC. roseoumCLS 104.52MK26827Present studyChina, YunnanC. roseoumCBS 104.52MK269228Present studyChina, YunnanC. roseoumCBS 104.52MK269228Present studyChina, YunnanC. chrysocrasKMRB 18041009MK120528[19]Korea, SeoulDarrymyces cyrtosporusPDD 17980NR_148190[28]USA, BethesdaD. flabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343523Present studyChina, YunnanD. confragosaCLZhao 4229MK343523Present studyChina, YunnanD. stereoidesCLZhao 4429MK343523Present studyChina, YunnanD. stereoidesCLZhao 3722KJ668554[19]Korea, SeoulD. stereoidesCLZhao 3722MK268896Present studyChina, ReijingEarliella scabrosaCui 6236KC867366[19]China, BeijingF albicansDai 10595NR, 153980[31]China, BeijingF lavodon ambrosiusCLZhao 3341MH14652Present studyChina, YunnanF. albicansDai 10595NR, 153980[31]China, KunnanF. albicansDai 10595NR, 153980[31]	C nulcherrimus	CLZbao 4379	MK343524	Present study	China Yunnan		
C. roseocarneumCLZhao 3562MK268877Present studyChina, YunnanC. roseumCBS 104.52MH856944[20]Netherlands, UpsalalaanCrustodontia chrysocreasCLZhao 4143MK269228Present studyChina, YunnanC. chrysocreasKMRB 18041009MK120528[19]Korea, SeoulDacrymyces cyrtosporusPDD 107980NR_148190[28]USA, BethesdaD. flabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MNS95677[30]Sweden, UppsalaDaconfragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCLZhao 4229MK343523Present studyChina, YunnanD. mollisKUC 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCui 283XU900526[19]China, YunnanD. stereoidesCui 283XU559270[19]China, YunnanEarliella scabrosaCui 6236KC867366[19]China, BeijingEarliella scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 3341MH114652Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, SunnanF. albicansCLZhao 4331MK343528Present studyChina, YunnanF. albicansCLZhao 4331MK343528Present studyChina, YunnanF. albicansCLZhao 4351MK343538Present studyChina	Corticium roseocarneum	CBS 330.66	MH858810	[20]	Netherlands Upsalalaan		
C. roseumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. roseumCBS 104.52MH856944[20]Netherlands, UpsalalaanC. chrysocreasCLZhao 4143MK20228Present studyChina, YunnanC. chrysocreasKMRB 18041009MK120528[19]Korea, SeoulDacrymyces cyrtosporusPDD 176696NR_148190[28]USA, BethesdaD. flabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN895677[30]Sweden, UppsalaDacdopsis confragosaCLZhao 4260MK343523Present studyChina, YunnanD. confragosaCui 2883KU900526[19]China, YunnanD. stergoidesCLZhao 4229MK343523Present studyChina, YunnanD. stergoidesCLZhao 4260MK943523Present studyChina, YunnanD. stergoidesCLZhao 4429MK343523Present studyChina, YunnanD. stergoidesCLZhao 3937MT19339Present studyChina, YunnanD. stergoidesCLZhao 3722MK268896Present studyChina, BeijingEarliella scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4311MH114652Present studyChina, WunnanFalbicansCLZhao 4331MK343526Present studyChina, WunnanF. anbrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK3	C roseocarneum	CLZbao 3562	MK268877	Present study	China Yunnan		
Crustodonita chrysocreasCLZhao 4143MR269228Present studyChina, YunnanC. chrysocreasKMRB 18041009MK120528[19]Korea, SeoulDacrymyces cyrtosporusPDD 107980NR_148190[28]USA, BethesdaD. flabeliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCLZhao 4429MK343523Present studyChina, WunnanD. confragosaCLZhao 4429MK343523Present studyChina, YunnanD. storonia mollisKUZ 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCLZhao 3722MK268896Present studyChina, YunnanE artiella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE scabrosaCLZhao 4416MK343526Present studyChina, YunnanF abicansDai 10595NR_153980[31]China, BeijingF labocansDai 10595NR_15328Present studyChina, YunnanF ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCLZhao 4351MK343538Present studyChina, YunnanGaioderma australeCLZhao	C roseum	CBS 104 52	MH856944	[20]	Netherlands Upsalalaan		
C. chrysocreasKMRB 1804100MK12052bTresht studyChina HanniDacrymyces cyrtosporusPDD 107980NR_148190[28]USA, BethesdaD. flabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCui 2883KU900526[19]China, BeijingDatonia mollisCLZhao 4429MK343523Present studyChina, YunnanD. storiagosaCLZhao 4429MK343533Present studyChina, YunnanD. storeoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCLZhao 3722MK268896Present studyChina, WunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, YunnanF. anbrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCui 8020JX290073[19]China, BeijingGaioderma australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, Quence, Quence	Crustodontia chrusocreas	CI Zbao 4143	MK269228	Present study	China Yunnan		
Dacrymyces cyrtosporusPDD 107980NR_148190[28]USA, BethesdaD. flabelliformisPDD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MNS95677[30]Sweden, UppsalaDacadaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCui 2883KU900526[19]China, YunnanD. confragosaCui 2883KU900526[19]Korea, SeoulD. mollisKU2 0121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCLZhao 4416MK343526Present studyChina, BeijingFibroporia albicansCLZhao 3341MH114652Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomentariusCui 8020JX290073[19]China, YunnanGanoderma australeCLZhao 4351MK343528Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGanoderma australeCLZhao 4351MK343528Present study	C chrusocreas	KMRB 18041009	MK120528	[19]	Korea Seoul		
DefibelijornisPDD 76696NR_166790[29]USA, BethesdaD. fibbelijornisPD 76696NR_166790[29]USA, BethesdaD. stillatusUPS F-939814MN595677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCui 2883KU900526[19]China, BeijingDatronia mollisCLZhao 4429MK343523Present studyChina, YunnanD. mollisKUC 20121123-22K[668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, BeijingE. scabrosaCLZhao 4416MK343526Present studyChina, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. sustraleHMAS 86595AY84184[19]China, BeijingGaodorina gunnanensisCLZhao 10504MN908253Present studyChina, YunnanG. sustraleHMAS 86595AY84184 <t< td=""><td>Dacrumuces curtosporus</td><td>PDD 107980</td><td>NR 148190</td><td>[28]</td><td>USA Bethesda</td></t<>	Dacrumuces curtosporus	PDD 107980	NR 148190	[28]	USA Bethesda		
D. JactillatisUPS F-939814MNS95677[30]Sweden, UppsalaDaedaleopsis confragosaCLZhao 4260MK343522Present studyChina, YunnanD. confragosaCui 2883KU900526[19]China, BeijingDatronia mollisCLZhao 4429MK343523Present studyChina, YunnanD. mollisKUC 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCLZhao 3722MK268896Present studyChina, YunnanEarliella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCui 8132JX559270[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, YunnanF. albicansDai 10595NR_153980[31]China, YunnanF. fomentariusCLZhao 3341MH114652Present studyChina, YunnanF. fomentariusCLZhao 4393MK343528Present studyChina, YunnanGauoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia gunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D flabelliformis	PDD 76696	NR 166790	[20]	USA Bethesda		
DaedaleopsisCl 2 ho 93014MR 95077[101]Swederl, OppsalaDaedaleopsisCl Zhao 4260MK 843522Present studyChina, BeijingD. confragosaCui 2883KU 900526[19]China, BeijingDatronia mollisCl Zhao 4429MK 343523Present studyChina, YunnanD. mollisKUC 20121123-22KJ 668554[19]Korea, SeoulD. stereoidesCl Zhao 9937MT 1935399Present studyChina, YunnanD. stereoidesCui 8132JX 559270[19]China, BeijingEarliella scabrosaCL Zhao 3722MK 268896Present studyChina, YunnanE. scabrosaCui 6236KC 867366[19]China, BeijingFibroporia albicansCl Zhao 3411MK 114652Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCl Zhao 3341MH 114652Present studyChina, YunnanF. ambrosiusHulcr 12546L C216231[32]USA, GainesvilleFomes fomentariusCui 8020JX 290073[19]China, BeijingGanoderma australeCl Zhao 4351MK 34358Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC 581962[33]Sweden, GoteborgGloeodontia yunnanensisCL Zhao 10504MN 908253Present studyChina, YunnanG. yunnanensisCL Zhao 11058M	D. stillatus	LIPS F-03081/	MNI595677	[20]	Swodon Unnsala		
D. confragosaCLZhao 4200MRC+5022Present studyChina, FurnariD. confragosaCui 2883KU900526[19]China, BeijingDatronia mollisCLZhao 4429MK343523Present studyChina, YunnanD. mollisKUC 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, WunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, BeijingEarliella scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingFilovodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHuler 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, WunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D. siliulus Daedaleoneis confragosa	$CI Z_{hao} 4260$	MK343522	[50] Procent study	China Yunnan		
D. tonji ugostiCui 1200ICS 3020[17]Cuinta, DeningDatronia mollisCLZhao 4429MK343523Present studyChina, YunnanD. mollisKUC 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, BeijingEarliella scabrosaCLZhao 4416MK343526Present studyChina, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, BeijingFlavodon ambrosiusCLZhao 4393MK343528Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanGanoderma australeCLZhao 4351MK343538Present studyChina, WunnanGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Ducunicopsis confragosa	C11i 2883	KI 1900526	[10]	China Boijing		
Dational monthsCLZ halo 4429MK845323Present studyClinia, HuntanD. mollisKUC 20121123-22KJ668554[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, WunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulc 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCLZhao 4351MK343538Present studyChina, WunnanGanoderma australeCLZhao 4351MK343538Present studyChina, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908253Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D. confrugosu Datronia mollia	CL Zbao 4429	MV242522	[17] Procent study	China, Deljilig		
D. montsKOC 20121125-22KJ660504[19]Korea, SeoulD. stereoidesCLZhao 9937MT193539Present studyChina, YunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingFiabicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCLZhao 4351MK343538Present studyChina, YunnanGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Dutroniu motifs	CLZIId0 4429	VI669EE4	[10]	Varia Sacul		
D. stereoidesCLZhao 9937M1193339Present studyChina, FunnanD. stereoidesCui 8132JX559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, WunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomentariusCLZhao 4393MK343528Present studyChina, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D. mouns	CL 7b == 0027	NJ008334	[19] Dura ana tanta dari	China Verman		
D. stereoidesCui 8132JA559270[19]China, BeijingEarliella scabrosaCLZhao 3722MK268896Present studyChina, YunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, BeijingF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomentariusCLZhao 4393MK343528Present studyChina, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, PunnanG. australeHMAS 86595AY884184[19]China, BeijingGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D. stereoides	CLZnao 9937	IVEE0270	Present study	China, Yunnan		
Earlielia scabrosaCLZhao 3722MR268896Present studyChina, FunnanE. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, PunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	D. stereotues	CL 7b = 2722	JA559270	[19] Dura ana tanta dari	China, beijing		
E. scabrosaCui 6236KC867366[19]China, BeijingFibroporia albicansCLZhao 4416MK343526Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan			WIK208896	Present study	China, Yunnan		
Fibroporia albicansCLZhao 4416MR343526Present studyChina, YunnanF. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	E. scabrosa	Cui 6236	KC867366		China, Beijing		
F. albicansDai 10595NR_153980[31]China, BeijingFlavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Fibroporia albicans	CLZhao 4416	MK343526	Present study	China, Yunnan		
Flavodon ambrosiusCLZhao 3341MH114652Present studyChina, YunnanF. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, BeijingG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	F. albicans	Dai 10595	NR_153980	[31]	China, Beijing		
F. ambrosiusHulcr 12546LC216231[32]USA, GainesvilleFomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Flavodon ambrosius	CLZhao 3341	MH114652	Present study	China, Yunnan		
Fomes fomentariusCLZhao 4393MK343528Present studyChina, YunnanF. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	F. ambrosius	Hulcr 12546	LC216231	[32]	USA, Gainesville		
F. fomentariusCui 8020JX290073[19]China, BeijingGanoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Fomes fomentarius	CLZhao 4393	MK343528	Present study	China, Yunnan		
Ganoderma australeCLZhao 4351MK343538Present studyChina, YunnanG. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	F. fomentarius	Cui 8020	JX290073	[19]	China, Beijing		
G. australeHMAS 86595AY884184[19]China, BeijingGeastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Ganoderma australe	CLZhao 4351	MK343538	Present study	China, Yunnan		
Geastrum pectinatumMJ 9030KC581962[33]Sweden, GoteborgGloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	G. australe	HMAS 86595	AY884184	[19]	China, Beijing		
Gloeodontia yunnanensisCLZhao 10504MN908252Present studyChina, YunnanG. yunnanensisCLZhao 11058MN908253Present studyChina, Yunnan	Geastrum pectinatum	MJ 9030	KC581962	[33]	Sweden, Goteborg		
G. yunnanensis CLZhao 11058 MN908253 Present study China, Yunnan	Gloeodontia yunnanensis	CLZhao 10504	MN908252	Present study	China, Yunnan		
	G. yunnanensis	CLZhao 11058	MN908253	Present study	China, Yunnan		

Table 1. List of species, specimens, and GenBank accession numbers of sequences used in this study.

# Table 1. Cont.

		GenBank Accession No.				
Species Name	Specimen No.	ITS	References	Country		
Gloeophyllum sepiarium	CBS 293.51	MH856865	[20]	Netherlands, Upsalalaan		
G. sepiarium	CLZhao 3473	MK268934	Present study	China, Yunnan		
Gomphus clavatus	EL 64/03	EU118628	[34]	Sweden, Goteborg		
Heterobasidion insulare	CLZhao 4071	MK268945	Present study	China, Yunnan		
H. insulare	Dai 13933	MT146489	[35]	China, Beijing		
Hexagonia glabra	CLZhao 3589	MK268955	Present study	China, Yunnan		
H. glabra	Cui 16796	MK192431	[36]	China, Beijing		
H. tenuis	CLZhao 3485	MK268956	Present study	China, Yunnan		
H. tenuis	Cui 8468	JX559277	[37]	China, Beijing		
Hymenochaete innexa	CLZhao 4433	MK343546	Present study	China, Yunnan		
H. innexa	001 855	M1596387	[19]	Italy, Iorino		
H. xerantica	CLZnao 2/89	MT24(0(F	Unpublished	China, Yunnan		
Hypnoaerma moniliforme	W11 0211 46	W11246965	Present study	China, Tunnan China, Taiwan		
н. monilijorme Н mutatum	CBS 201 36	NC926264 MH855803	[30]	Nothorlands Unsalalaan		
H mutatum	CI 7bao 10019	MT246969	[20] Present study	China Vunnan		
H nudicenhalum	CBS 125855	MH863811	[20]	Netherlands Upsalalaan		
H nudicephalum	CL Zhao 3481	MK269020	Present study	China. Yunnan		
H. transiens	CLZhao 9628	MT246975	Present study	China, Yunnan		
H. transiens	NH 12304	DO677504	[39]	Sweden Goteborg		
Hysterangium atlanticum	Fungos 2115	NR 165211	[40]	Slovenia, Liubliana		
Irvex lacteus	CLZhao 4338	MK343588	Present study	China, Yunnan		
I. lacteus	FD-9	KP135026	[25]	USA, Worcester		
Junghuhnia crustacea	CLZhao 3588	MK269088	Present study	China, Yunnan		
J. crustacea	X 1127	JN710554	[41]	Finland, Helsinki		
J. nitida	CLZhao 3474	MK269086	Present study	China, Yunnan		
J. nitida	KHL 11903	EU118638	[34]	Sweden, Goteborg		
Laxitextum incrustatum	CLZhao 9273	MT159974	Present study	China, Yunnan		
L. incrustatum	STMA 14285	KT722621	[42]	Germany, Braunschweig		
Lenzites betulinus	CLZhao 10151	MT193546	Present study	China, Yunnan		
L. betulinus	Cui 7095	JX290075	[19]	China, Beijing		
Lopharia mirabilis	CLZhao 3518	MK269105	Present study	China, Yunnan		
L. mirabilis	Dai 13722	MF626346	[43]	China, Beijing		
Lyomyces bambusinus	CLZhao 3675	MN945969	Present study	China, Yunnan		
L. bambusinus	CLZhao 4840	MN945971	Present study	China, Yunnan		
L. cremeus	CLZhao 4138	MIN945974	Present study	China, Yunnan		
L. cremeus	CLZnao 8295 CLZhao 4516	MIN945972	Present study	China, Yunnan		
L. macrosporus	CLZhao 4516	MN1945977	Present study	China, Tunnan		
L. mucrosporus	CLZhao 4475	MN045083	Prosont study	China, Yunnan		
L. wulliangshanensisd	CI Zhao 4108	MN945980	Present study	China Yunnan		
Megasnoria ellinsoidea	CLZhao 10360	MT193554	Present study	China Yunnan		
M. ellipsoidea	Cui 5222	IO314367	[19]	China, Beijing		
M. major	CLZhao 10363	MT193555	Present study	China, Yunnan		
M. major	Yuan 1183	IO314365	[19]	China, Beijing		
Megasporoporiella		, <u>2</u> ,	[]			
pseudocavernulosa	CLZhao 9339	MK894044	Present study	China, Yunnan		
M. pseudocavernulosa	Cui 11106	KX900657	[19]	China, Beijing		
M. subcavernulosa	CLZhao 4442	MK343599	Present study	China, Yunnan		
M. subcavernulosa	Cui 10050	JQ314357	[19]	China, Beijing		
Microporus vernicipes	CLZhao 3672	MK269132	Present study	China, Yunnan		
M. vernicipes	Dai 7252	KX880619	[19]	China, Beijing		
M. xanthopus	CLZhao 4280	MT159985	Present study	China, Yunnan		
M. xanthopus	Dai 12076	KX880620	[19]	China, Beijing		
Neodatronia sinensis	CLZhao 9471	MT193563	Present study	China, Yunnan		
N. sinensis	Cui 9949	KX900663	[19]	China, Beijing		
Oxychaete cervinogilva	CLZhao 3462	MK269143	Present study	China, Yunnan		
O. cervinogilva	Schigel 5216	KX752596	[44]	Finland, Helsinki		
Peniopnora cinerea	0911 1E52/A5	LIN808982	[19]	Spain, Sevilla		
P. cinerea	CLZnao 9254	MI 159975	Present study	China, Yunnan		
r. incurnuta D incarnata	CD5 450.72	IVIF100U218 MT150074	[40] Procent study	China Vunnan		
F. Incurnunu P nesudopersicolor	CLZ11d0 9009 CL 7hao 9415	MT150077	Present study	China Vunnan		
F. pesudopersicolor D negudopersicolor	CBS 229 44	MHQEQQ11	[20]	Nothorlanda Unsalalaan		
1. pseudoversicoior Penionhorella fissurata	CL7hao 11412	MNI864261	[40] Present study	China Vunnan		
P. fissurata	CLZhao 4539	MN864259	Present study	China, Yunnan		
P. pubera	CLZhao 9336	MK894055	Present study	China, Yunnan		

 Table 1. Cont.

GenBank Accession No.							
Species Name	Specimen No.	ITS	References	Country			
P. pubera	NH 10380	DQ647504	[45]	Sweden, Goteborg			
P. rude	CLZhao 3360	MH114830	Present study	China, Yunnan			
P. rude	Wu 9307-39	DQ647499	[45]	Sweden, Goteborg			
Perenniporia luteola	CLZhao 10334	MT193564	Present study	China, Yunnan			
P. luteola	Dai 13094	KX900679	[19]	China, Beijing			
P. mopanshanensis	CLZhao 4677	MK404422	Present study	China, Yunnan			
P. mopanshanensis	CLZhao 5145	MH784912	Present study	China, Yunnan			
Phaeophlebiopsis peniophoroides	CLZhao 3637	MK269159	Present study	China, Yunnan			
P. peniophoroides	FP-150577	KP135417	[25]	USA, Worcester			
Phallus impudicus	KHTW 08001	MG678529	[46]	Brazil, Manaus			
Phanerochaete	CLZhao 9280	MT177305	Present study	China, Yunnan			
D angustocustidiata	KUC20121102-15	K1668402	[10]	Koroa Socul			
D australic	CL Zhao 3571	MK269240	[17] Procent study	China Vunnan			
D australia	ULIR 7105 Sp	VD125081	[25]	USA Worcostor			
T. UUSIIUIIS	CL Zhao 4267	NI 135081 MI/ 404420	[25] Procent study	China Vunnan			
P. concrescens	CLZIIdO 4307	WIN404450	Fresent study	Chilla, Tulhan Bussis, Ssint Datarahura			
P. concrescens	CL Zhao 4447	NF 994552 NF 242612	[47] Drecent study	China Vunnan			
P. sunguineocurnosu		WIN545015	Fresent study				
P. sanguineocarnosa	HHB-2189	KP135124	[25]	USA, worcester			
P. soraiaa	CLZnao 340/	MIK269174	Present study	China, Yunnan			
P. soraida	FD-241	KP135136	[25]	USA, Worcester			
Phlebiopsis castanea	CLZhao 3501	MK269230	Present study	China, Yunnan			
P. castanea	Spirin 5295	KX752610	[44]	Finland, Helsinki			
P. crassa	CLZhao 3339	MT177325	Present study	China, Yunnan			
P. crassa	KKN-86	KP135394	[25]	USA, Worcester			
khomboidia wuliangshanensis	CLZhao 4406	MK860715	Present study	China, Yunnan			
R. wuliangshanensis	CLZhao 4411	MK860716	Present study	China, Yunnan			
Rigidoporus pouzarii	CLZhao 4524	MK404499	Present study	China, Yunnan			
R. pouzarii	MJ 123_93	JQ409464	[19]	Czech Republic, South Bohemia			
Russula recondita	LUG 19058	NR_147635	[48]	USA, Bethesda			
Sebacina incrustans	JMP 0084	EU819442	[49]	USA, Wisconsin			
Serpula dendrocalami	CLZhao 3136	MK863399	Present study	China, Yunnan			
, S. dendrocalami	CLZhao 3607	MK863404	Present study	China, Yunnan			
Sistotrema brinkmannii	ATCC 36164	DQ899095	[19]	USA, Stillwater			
S. brinkmannii	CLZhao 9324	MT159987	Present study	China, Yunnan			
Sistotremastrum		NUM4000/1	[10]				
guttuliferum	DHI-F/35a	11111488061	[19]	USA, west Larayette			
S. guttuliferum	CLZhao 9771	MT159990	Present study	China, Yunnan			
Steccherinum bourdotii	CLZhao 4297	MK343648	Present study	China, Yunnan			
S. bourdotii	U 107	MN749593	[50]	Arabia, Hail			
S. ochraceum	CLZhao 4154	MK404504	Present study	China, Yunnan			
S. ochraceum	LE-BIN 3984	MK795064	[19]	Russia, Saint Petersburg			
Terana coerulea	CLZhao 9803	MT177329	Present study	China, Yunnan			
T. coerulea	FP-104073	KP134980	[25]	USA, Worcester			
Thelephora terrestris	JMP 0086	EU819444	[49]	USA, La Crosse			
Tinctoporellus epimiltinus	CLZhao 3605	MK269225	Present study	China, Yunnan			
T. epimiltinus	PR1147	KY948721	[51]	USA, Massachusetts			
Trametes ellipsosopia	CLZhao 3774	MK269226	Present study	China, Yunnan			
T. ellipsospora	Cui 6259	JN048767	[52]	China, Beijing			
T. hirsuta	CLZhao 3572	MK269220	Present study	China, Yunnan			
T. hirsuta	TH 158	MF960866	[53]	Czech Republic, Brno			
T. pavonia	CLZhao 3740	MK269222	Present study	China, Yunnan			
T. pavonia	PR2178	JN164959	[54]	USA, Worcester			
T. versicolor	CLZhao 4170	MK269115	Present study	China, Yunnan			
T. versicolor	F-1599	MN749366	[19]	USA, West Lafavette			
Trametopsis cervina	CLZhao 9449	MW418032	Present study	China, Yunnan			
T. cervina	TJV 93 216T	JN165020	[54]	USA, Worcester			
Trechispora echinospora	MA-Fungi 82485	JX392847	[55]	Spain, Madrid			
T. nivea	A 1683	MT458536	[19]	Greece, Athens			
T. nivea	CLZhao 10154	MT159996	Present study	China, Yunnan			
Truncospora ochroleuca	CLZhao 4350	MK343685	Present study	China, Yunnan			
T. ochroleuca	BCC 16648	KP059107	[19]	Thailand, Khlong			

Sequencher 4.6 (GeneCodes, Ann Arbor, MI, USA) was used to edit the DNA sequence. Sequences were aligned in MAFFT 7 (http://mafft.cbrc.jp/alignment/server/, accessed on 15 November 2021) using the "G-INS-i" strategy and manually adjusted in BioEdit [56]. Sequence of Dacrymyces from type species as *D. stillatus* Nees obtained from GenBank was used as an outgroup to root tree following James et al. [1] in ITS analysis (Figure 1). Sequences from type materials of *Dacrymyces flabelliformis* Burds. and Laursen and *D. cyrtosporus* Shirouzu acquired from GenBank were utilized as another outgroup to root tree following James et al. [1] in ITS analysis (Figure 2).

Maximum parsimony analysis was applied to the ITS dataset sequences. Approaches to phylogenetic analysis followed Zhao and Wu [57] and the tree construction procedure was performed in PAUP\* version 4.0b10 [58]. All characters were equally weighted and gaps were treated as missing data. Trees were inferred using the heuristic search option with TBR branch swapping and 1000 random sequence additions. Max-trees were set to 5000, branches of zero length were collapsed and all parsimonious trees were saved. Clade robustness was assessed using a bootstrap (BT) analysis with 1000 replicates [59]. Descriptive tree statistics tree length (TL), consistency index (CI), retention index (RI), rescaled consistency index (RC), and homoplasy index (HI) were calculated for each Maximum Parsimonious Tree (MPT) generated. Sequences were also analyzed using maximum likelihood (ML) with RAxML-HPC2 through the Cipres Science Gateway [60]. Branch support (BS) for ML analysis was determined by 1000 bootstrap replicates.

MrModeltest 2.3 [61] was used to determine the best-fit evolution model for each data set for Bayesian inference (BI). Bayesian inference was calculated with MrBayes3.1.2 with a general time reversible (GTR+I+G) model of DNA substitution and a gamma distribution rate variation across sites [62]. Four Markov chains were run for 2 runs from random starting trees for 760 thousand generations (Figure 1) and 6.4 million generations (Figure 2), and trees were sampled every 100 generations. The first one-fourth generations were discarded as burn-in. A majority rule consensus tree of all remaining trees was calculated. Branches were considered as significantly supported if they received maximum likelihood bootstrap (BS) >70%, maximum parsimony bootstrap (BT) >50%, or Bayesian posterior probabilities (BPP) >0.95.



**Figure 1.** Maximum parsimony strict consensus tree illustrating the phylogeny of 9 orders in Agaricomycetes based on ITS sequences. Branches are labeled with maximum likelihood bootstrap >70%, parsimony bootstrap >50% and Bayesian posterior probabilities >0.97, respectively. The 9 orders conducted in present studies are labeled in light yellow.



Figure 2. Cont.



**Figure 2.** Maximum parsimony strict consensus tree illustrating the phylogeny of 85 species in Agaricomycetes based on ITS sequences. Branches are labeled with maximum likelihood bootstrap >70%, parsimony bootstrap >50% and Bayesian posterior probabilities >0.97, respectively. The sequences of collections labeled in orange are downloaded from GenBank, while those in black are generated in this study.

#### 3. Results

#### 3.1. Molecular Phylogeny

The ITS dataset (Figure 1) included sequences from 18 fungal specimens representing 18 species. The dataset had an aligned length of 1017 characters, of which 326 characters were constant, 272 parsimony-uninformative, and 419 parsimony-informative. Maximum parsimony analysis yielded three equally parsimonious trees (TL = 2801, CI = 0.4763, HI = 0.5237, RI = 0.0259, RC = 0.0123). The best-fit model for ITS alignment estimated and applied in the Bayesian method was GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet (1,1,1,1). The Bayesian method resulted in a similar topology with an average standard deviation of split frequencies = 0.009891 (BI).

The phylogeny (Figure 1) inferred from ITS sequences demonstrated that ninetyfive species nested in nine orders, including Auriculariales, Boletales, Cantharellales, Corticiales, Gloeophyllales, Hymenochaetales, Polyporales, Russulales and Trechisporales of Agaricomycetes.

The ITS dataset (Figure 2) included sequences from 172 fungal specimens representing 87 species. The dataset had an aligned length of 1107 characters, of which 351 characters were constant, 52 parsimony-uninformative, and 704 parsimony-informative. Maximum parsimony analysis yielded 10 equally parsimonious trees (TL = 7174, CI = 0.2319, HI = 0.7681, RI = 0.6952, RC = 0.1613). The best-fit model for ITS alignment estimated and applied in the Bayesian method was GTR+I+G, lset nst = 6, rates = invgamma; prset statefreqpr = dirichlet (1,1,1,1). The Bayesian method resulted in a similar topology with an average standard deviation of split frequencies = 0.012495 (BI), and the effective sample size (ESS) across the two runs is the double of the average ESS (avg ESS) = 223.5.

The phylogeny (Figure 2) inferred from ITS sequences demonstrated that 2454 specimens of wood-decaying fungi belonged to 85 taxa and all of them nested into class Agaricomycetes.

#### 3.2. Checklist

An alphabetical list (according to genus name) of wood-decaying fungi identified in these investigations is given below. The authors of scientific names are according to the second edition of *Authors of Fungal Names* (http://www.indexfungorum.org/AuthorsO fFungalNames.htm, accessed on 15 November 2021). Substrate and collecting data are provided after the name of each species. The hosts are listed alphabetically, and within the same host tree, they are arranged by the following order: living tree, dead standing tree, trunk, fallen branch and stump. The collectors and collection numbers are listed alphabetically as well [6,63].

*Abundisporus quercicola* Y.C. Dai, in Dai, Niemelä and Kinnunen, Ann. bot. fenn. 39(3): 171 (2002).

Specimens examined: on dead tree of angiosperm, 9 January 2019, CLZhao 10238; on the trunk of angiosperm, 10 January 2019, CLZhao 10,381 (SWFC).

Distribution: Asia—China [64].

Aleurodiscus botryosus Burt, Ann. Mo. bot. Gdn 5: 198 (1918).

Specimens examined: on the fallen angiosperm branch, 5 October 2017, CLZhao 4333 (SWFC). Distribution: Asia—China; Europe—France Portugal, Spain the United Kingdom;

North America—Canada the United States; Oceania—Australia, New Zealand [64]. Aleurodiscus cerussatus (Bres.) Höhn. and Litsch., Sber. Akad. Wiss. Wien, Math.-

naturw. Kl., Abt. 1 116: 808 (1907).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4715 (SWFC).

Distribution: Asia—China; Europe—Denmark, Estonia, Italy, Norway, Russia, Spain, Sweden; North America—the United States; South America—Argentina; Oceania—Australia [64].

*Aleurodiscus isabellinus* S.H. He and Y.C. Dai, in Tian, Ghobad-Nejhad, He and Dai, MycoKeys 37: 100 (2018).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4102; 6 October 2017, CLZhao 4405, CLZhao 4600; 4 January 2019, CLZhao 9300; on the fallen branch of angiosperm, 5 October 2017, CLZhao 4369, CLZhao 4435, CLZhao 4510, CLZhao 4564, CLZhao 4718; 4 January 2019, CLZhao 9284, CLZhao 9359, CLZhao 9364, CLZhao 9615, CLZhao 9643; 7 January 2019, CLZhao 9781, CLZhao 9861; 9 January 2019, CLZhao 10,095 (SWFC).

Distribution: Asia—China [64].

*Aleurodiscus mirabilis* (Berk. and M.A. Curtis) Höhn., Sber. Akad. Wiss. Wien, Math.-naturw. Kl., Abt. 1 118: 818 (1909).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4621 (SWFC).

Distribution: Africa—Gabon, Mauritius, Tanzania; Asia—China, Japan, Sri Lanka; North America—Cuba, Jamaica, the United States; South America—Brazil, Colombia, Ecuador, Venezuela; Oceania—Australia, New Zealand [64].

*Aleurodiscus subroseus* S.H. He and Y.C. Dai, in Tian, Ghobad-Nejhad, He and Dai, MycoKeys 37: 102 (2018).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4722 (SWFC).

Distribution: Asia—China [64].

Asterostroma muscicola (Berk. and M.A. Curtis) Massee, J. Linn. Soc., Bot. 25(no. 170): 155 (1889).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3577 (SWFC).

Distribution: Africa—Malawi, Rwanda, Tanzania; Asia—China, India, Indonesia; North America—Bahamas, Cuba, Canada, Jamaica, Mexico, the United States; South America—Brazil, Colombia, Venezuela; Oceania—Australia [64].

*Auricularia asiatica* Bandara and K.D. Hyde, in Bandara, Karunarathna, Phillips, Mortimer, Xu, Kakumyan and Hyde, Phytotaxa 292(1): 24 (2016).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3652 (SWFC). Distribution: Asia—China, Thailand [64].

*Auricularia cornea* Ehrenb., in Nees von Esenbeck (Ed.), Horae Phys. Berol.: 91 (1820). Specimens examined: on the dead tree of angiosperm, 9 January 2019, CLZhao 10129; on the trunk of angiosperm, 5 October 2017, CLZhao 4245, CLZhao 4282, CLZhao 4306, CLZhao 4483; 4 January 2019, CLZhao 9272, CLZhao 9330; 9 January 2019, CLZhao 10009, CLZhao 10,324 (SWFC).

Distribution: Africa—South Africa, Tanzania; Asia—China, Indonesia; North America— Mexico, the United States; South America—Brazil [64].

Auricularia delicata (Mont. ex Fr.) Henn., Bot. Jb. 17: 492 (1893).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4403, CLZhao 4582 (SWFC).

Distribution: Asia—China, India; North America—Mexico, the United States [64]. *Bjerkandera adusta* (Willd.) P. Karst., Meddn Soc. Fauna Flora fenn. 5: 38 (1879).

Specimens examined: on the fallen branch of angiosperm, 5 October 2017, CLZhao 4310; 6 October 2017, CLZhao 4371, CLZhao 4374, CLZhao 4376, CLZhao 4427, CLZhao 4444, CLZhao 4546, CLZhao 4625, CLZhao 4637; 7 January 2019, CLZhao 9844; 4 January 2019, CLZhao 9345; 5 January 2019, CLZhao 9508; 10 January 2019, CLZhao 10385, CLZhao 10,404 (SWFC).

Distribution: Asia—China, India, Japan, Pakistan, Thailand, Vietnam; Europe—Bulgaria, Finland, Germany, Italy, Norway, Poland, Russia, Spain, Sweden, the United Kingdom, Ukraine; North America—Canada, the United States [64].

*Byssomerulius corium* (Pers.) Parmasto, Eesti NSV Tead. Akad. Toim., Biol. seer 16(4): 383 (1967).

Specimens examined: on the trunk of angiosperm, 4 January 2019, CLZhao 9318; on the fallen branch of angiosperm, 4 January 2019, CLZhao 9390, CLZhao 10,110 (SWFC).

Distribution: Asia—China, Turkey; Europe—Belgium, Croatia, Estonia, Finland, France, Germany, Greece, Ireland, Italy, Norway, Poland, Portugal, Romania, Serbia, Spain, Sweden, Switzerland, the United Kingdom, Ukraine [64].

*Ceriporia lacerata* N. Maek., Suhara and R. Kondo, in Suhara, Maekawa, Kaneko, Hattori, Sakai and Kondo, Mycotaxon 86: 342 (2003).

Specimens examined: on the trunk of angiosperm, 3 October 2017, CLZhao 3693 (SWFC).

Distribution: Asia—China, Japan; Europe—Finland; North America—the United States [64].

Cerrena albocinnamomea (Y.C. Dai and Niemelä) H.S. Yuan, Mycol. Progr. 13(2): 362 (2013).

Specimens examined: on the fallen branch of *Alnus nepalensis*, 4 January 2019, CLZhao 9281; on the trunk of angiosperm, 5 October 2017, CLZhao 4197; 4 January 2019, CLZhao 9270, CLZhao 9342; on the fallen branch of angiosperm, 5 October 2017, CLZhao 4193 (SWFC).

Distribution: Asia—China, India, Japan, South Korea [64].

Cerrena zonata (Berk.) H.S. Yuan, Mycol. Progr. 13(2): 363 (2013).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3581; 5 October 2017, CLZhao 4178; 6 January 2019, CLZhao 9633, CLZhao 9641; on the fallen branch of angiosperm, 2 October 2017, CLZhao 3502 (SWFC).

Distribution: Asia—China, India, Japan, Thailand, Vietnam; Europe—Russia; South America—Argentina; Oceania—Australia, New Zealand [64].

*Climacodon pulcherrimus* (Berk. and M.A. Curtis) Nikol., Flora Plantarum Cryptogamarum URSS 6, Fungi 6(Fungi, 2): 194 (1961).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4379 (SWFC).

Distribution: Asia—China; Europe—Belarus, France, Spain, Ukraine [64].

Corticium roseocarneum (Schwein.) Hjortstam, Windahlia 23: 2 (1998).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4719; on the fallen branch of angiosperm, 6 October 2017, CLZhao 4545, CLZhao 4735; 7 January 2019, CLZhao 9815; on the trunk of *Pinus*, 2 October 2017, CLZhao 3562 (SWFC).

Distribution: Asia—China, Japan; Europe—France, Norway; North America—Canada [64]. *Crustodontia chrysocreas* (Berk. and M.A. Curtis) Hjortstam and Ryvarden, Syn. Fung. (Oslo) 20: 36 (2005).

Specimens examined: on the fallen branch of angiosperm, 5 October 2017, CLZhao 4143; 7 January 2019, CLZhao 9831 (SWFC).

Distribution: Asia—China; Europe—Belarus, France [64].

*Dacryobolus angiospermarum* S.H. He, in Xu, Liu, Wu and He, Phytotaxa 365(2): 190 (2018). Specimens examined: on the fallen branch of *Pinus*, 3 October 2017, CLZhao 3706 (SWFC). Distribution: Asia—China [64].

*Daedaleopsis confragosa* (Bolton) J. Schröt., in Cohn, Krypt.-Fl. Schlesien (Breslau) 3.1(25–32): 492 (1888).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4260; on the stump of angiosperm, 10 January 2019, CLZhao 10379, CLZhao 10,447 (SWFC).

Distribution: Africa—Nigeria, South Africa; Asia—China, Azerbaijan, Bhutan, India, Iran, Japan, Malaysia, Nepal, North Korea; Europe—Andorra, Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, Finland, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Netherlands, Norway, Poland, Romania, Russia, Serbia, Slovakia, Slovenia, Spain, Sweden, Switzerland, the United Kingdom, Ukraine; North America—Canada, Mexico, the United States; South America—Brazil, Venezuela; Oceania—Australia [64].

Datronia mollis (Sommerf.) Donk, Persoonia 4(3): 338 (1966).

Specimens examined: on dead tree of angiosperm, 6 October 2017, CLZhao 4429; on the trunk of angiosperm, 4 January 2019, CLZhao 9326 (SWFC).

Distribution: Asia—China; Europe—Russia; North America—the United States [64]. *Datronia stereoides* (Fr.) Ryvarden, Blyttia 25: 168 (1967).

Specimens examined: on the fallen branch of angiosperm, 7 January 2019, CLZhao 9937 (SWFC).

Distribution: Asia—China, Japan, Kyrgyzstan, North Korea; Europe—Austria, Finland, France, Hungary, Norway, Russia, Spain, Sweden, Switzerland, the United Kingdom; North America—Canada, Costa Rica, Honduras, Mexico, the United States; South America—Brazil, Colombia, Guyana, Paraguay, Venezuela; Oceania—Australia [64].

*Earliella scabrosa* (Pers.) Gilb. and Ryvarden, Mycotaxon 22(2): 364 (1985).

Specimens examined: on the trunk of angiosperm, 3 October 2017, CLZhao 3688, CLZhao 3701, CLZhao 3722, CLZhao 3754, CLZhao 3766; on the stump of angiosperm, 3 October 2017, CLZhao 3730 (SWFC).

Distribution: Asia—China, India, Malaysia, Philippines, Sri Lanka; North America—the United States [64].

*Fibroporia albicans* B.K. Cui and Yuan Y. Chen, in Chen, Li and Cui, Phytotaxa 203(1): 51 (2015).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4416 (SWFC). Distribution: Asia—China [64].

*Flavodon ambrosius* D.R. Simmons, You Li, C.C. Bateman and J. Hulcr, Mycotaxon 131(2): 279 (2016).

Specimens examined: on the trunk of angiosperm, 1 October 2017, CLZhao 3341 (SWFC). Distribution: Asia—China; North America—the United States [64].

*Fomes fomentarius* (L.) Fr., Summa veg. Scand., Sectio Post. (Stockholm): 321 (1849). Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4393 (SWFC).

Distribution: Asia—China, Japan; Europe—Austria, Belgium, Czech Republic, Denmark, Estonia, Finland, France, Germany, Italy, Luxembourg, Netherlands, Norway, Poland, Russia, Slovenia, Spain, Sweden, the United Kingdom, Ukraine; North America—Canada, the United States [64]. *Ganoderma australe* (Fr.) Pat., Bull. Soc. mycol. Fr. 5(2,3): 65 (1889) (Figure 3).



**Figure 3.** Basidiomata in situ of *Ganoderma australe*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = (**D**) = 2 mm.

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4351 (SWFC).

Distribution: Asia—China; Europe—France, the United Kingdom; North America—Canada, the United States; Oceania—Australia, New Zealand [64].

*Gloeodontia yunnanensis* C.L. Zhao, in Chen, Shi, Wu and Zhao, Phytotaxa 432(2): 115 (2020).

Specimens examined: on angiosperm trunk, 10 January 2019, CLZhao 10,504 (SWFC). Distribution: Asia—China [64].

*Gloeophyllum sepiarium* (Wulfen) P. Karst. (as 'Gleophyllum'), Bidr. Känn. Finl. Nat. Folk 37: 79 (1882) (Figure 4).



**Figure 4.** Basidiomata in situ of *Gloeophyllum sepiarium*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = (**D**) = 2 mm.

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3473, CLZhao 3500, CLZhao 3555, CLZhao 3559, CLZhao 3575; 5 October 2017, CLZhao 4155; 6 October 2017, CLZhao 4532; on the fallen branch of *Pinus*, 3 October 2017, CLZhao 3680 (SWFC).

Distribution: Asia—China, Japan; Europe—Russia; North America—the United States [64].

Gyrodontium sacchari (Spreng.) Hjortstam, Mycotaxon 54: 186 (1995).

Specimens examined: on the stump of angiosperm, 3 October 2017, CLZhao 3572 (SWFC). Distribution: Asia—China, India; Europe—Belgium, Italy, Sweden; North America—

## Mexico [64].

*Heterobasidion insulare* (Murrill) Ryvarden (as 'insularis'), Norw. Jl Bot. 19: 237 (1972) (Figure 5).



**Figure 5.** Basidiomata in situ of *Heterobasidion insulare*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = (**D**) = 1 mm.

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4071, CLZhao 4132, CLZhao 4145, CLZhao 4220, CLZhao 4363; on the stump of angiosperm, 5 October 2017, CLZhao 4224; on the trunk of *Pinus*, 5 October 2017, CLZhao 4216 (SWFC).

Distribution: Asia—China, India, Japan, Korea, Myanmar, Nepal, South Korea; Europe—Russia [64].

Hexagonia glabra (P. Beauv.) Ryvarden (as 'glaber'), Mycotaxon 72: 216 (1999).

Specimens examined: on the fallen branch of angiosperm, 2 October 2017, CLZhao 3589 (SWFC).

Distribution: Africa—Tanzania, Mauritius; Asia—China, India, Japan, Philippines, Thailand, Vietnam; North America—Costa Rica; South America—Brazil; Oceania—Australia [64].

*Hexagonia tenuis* (Fr.) Fr., Epicr. syst. mycol. (Upsaliae): 498 (1838). Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3485, CLZhao 3498 (SWFC).

Distribution: Asia—China, India, Japan, Malaysia, Philippines, Sri Lanka. Oceania—Australia [64].

*Hydnoporia corrugata* (Fr.) K.H. Larss. and Spirin, in Miettinen, Larsson and Spirin, Fungal Systematics and Evolution 4: 88 (2019).

Specimens examined: on the fallen branch of angiosperm, 7 January 2019, CLZhao 9913; 9 January 2019, CLZhao 10170, CLZhao 10207; 10 January 2019, CLZhao 10344, CLZhao 10,362 (SWFC).

Distribution: Asia—China, Japan, Sri Lanka; Europe—Denmark, France, Germany, Ireland, Netherlands, Norway, Russia, Spain, Sweden, Switzerland, the United Kingdom, Yugoslavia; North America—Canada, the United States; South America—Argentina, Colombia; Oceania—New Zealand [64].

Hymenochaete berteroi Pat., Bull. Soc. mycol. Fr. 10(2): 78 (1894).

Specimens examined: on the fallen branch of angiosperm, 5 October 2017, CLZhao 4203, CLZhao 4328 (SWFC).

Distribution: Africa—Kenya, Malawi, Tanzania; Asia—China; North America—Costa Rica; South America—Venezuela [64].

Hymenochaete innexa G. Cunn., Trans. Roy. Soc. N.Z. 85(1): 47 (1957).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4433, CLZhao 4458, CLZhao 4584, CLZhao 4586, CLZhao 4731; 3 January 2019, CLZhao 9255; 5 January 2019, CLZhao 9528 (SWFC).

Distribution: Africa—Seychelles; Asia—China, India; North America—Jamaica, the United States; South America—Brazil; Oceania—Australia, New Zealand [64].

*Hyphoderma moniliforme* (P.H.B. Talbot) Manjón, G. Moreno and Hjortstam, in Hjortstam, Manjón and Moreno, Mycotaxon 33: 261 (1988).

Specimens examined: on the fallen branch of angiosperm, 9 January 2019, CLZhao 10176, CLZhao 10232, CLZhao 10242; 10 January 2019, CLZhao 10,386 (SWFC).

Distribution: Asia—China; Europe—Spain [64].

Hyphoderma mutatum (Peck) Donk, Fungus, Wageningen 27: 15 (1957).

Specimens examined: on the fallen branch of angiosperm, 9 January 2019, CLZhao 10019, CLZhao 10050, CLZhao 10104, CLZhao 10139, CLZhao 10141, CLZhao 10,203 (SWFC).

Distribution: Asia—China, Japan; Europe—Denmark, Finland, Norway, Sweden; North America—Canada, the United States [64].

*Hyphoderma nudicephalum* Gilb. And M. Blackw., Mycotaxon 33: 378 (1988). Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3481 (SWFC). Distribution: Asia—China, Japan, Korea; Europe—Denmark, Spain; Oceania—Australia, New Zealand [64].

*Hyphoderma transiens* (Bres.) Parmasto, Consp. System. Corticiac. (Tartu): 114 (1968) Specimens examined: on the fallen branch of angiosperm, 6 January 2019, CLZhao 9628 (SWFC).

Distribution: Asia—China, Japan [64].

Irpex lacteus (Fr.) Fr., Elench. fung. (Greifswald) 1: 142 (1828) (Figure 6).



**Figure 6.** Basidiomata in situ of *Irpex lacteus*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = (**D**) = 1 mm.

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4338, CLZhao 4365; 6 October 2017, CLZhao 4579; 4 January 2019, CLZhao 9438; 6 January 2019, CLZhao 9765; 7 January 2019, CLZhao 9816, CLZhao 9824, CLZhao 9855, CLZhao 9883, CLZhao 9922; on the fallen branch of angiosperm, 6 October 2017, CLZhao 4609, CLZhao

4730, CLZhao 9398, CLZhao 9612, CLZhao 9728, CLZhao 9748; 7 January 2019, CLZhao 9878, CLZhao 9888, CLZhao 10,113 (SWFC).

Distribution: Asia—China, Japan; Europe—France, the United Kingdom; North America the United States [64].

Junghuhnia crustacea (Jungh.) Ryvarden, Persoonia 7(1): 18 (1972).

Specimens examined: on the fallen branch of angiosperm, 2 October 2017, CLZhao 3588 (SWFC).

Distribution: Asia—China; North America—the United States; South America—Brazil [64]. *Junghuhnia nitida* (Pers.) Ryvarden, Persoonia 7(1): 18 (1972).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3650; 6 October 2017, CLZhao 4439, CLZhao 4455, CLZhao 4631; on the fallen branch of angiosperm, 2 October 2017, CLZhao 3474, CLZhao 3480; 5 October 2017, CLZhao 4219; 6 October 2017, CLZhao 4702; 5 January 2019, CLZhao 9447, CLZhao 9588; 6 January 2019, CLZhao 9667, CLZhao 9707; 7 January 2019, CLZhao 9854, CLZhao 9884, CLZhao 9919, CLZhao 9943, CLZhao 9969, CLZhao 9998; 9 January 2019, CLZhao 10082, CLZhao 10097, CLZhao 10178, CLZhao 10186, CLZhao 10211, CLZhao 10252, CLZhao 10271, CLZhao 10323; 10 January 2019, CLZhao 10350, CLZhao 10436, CLZhao 10458, CLZhao 10476, CLZhao 10497, CLZhao 10,535 (SWFC).

Distribution: Asia—China, Japan; Europe—France, the United Kingdom; North America—the United States [64].

*Laxitextum incrustatum* Hjortstam and Ryvarden, Mycotaxon 13(1): 35 (1981).

Specimens examined: on the trunk of angiosperm, 4 January 2019, CLZhao 9273 (SWFC). Distribution: Africa—Tanzania, Ethiopia; Asia—China; North America—the United States; South America—Brazil [64].

*Lenzites betulinus* (L.) Fr. (as 'betulina'), Epicr. syst. mycol. (Upsaliae): 405 (1838). Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4268, CLZhao 4281; 9 January 2019, CLZhao 10151; on the stump of angiosperm, 10 January 2019, CLZhao 10425, CLZhao 10454, CLZhao 10498, CLZhao 10,516 (SWFC).

Distribution: Asia—China, Japan, North Korea, Philippines; Europe—Austria, Belgium, Denmark, Estonia, Finland, France, Germany, Netherlands, Norway, Russia, Slovenia, Spain, Sweden, the United Kingdom; North America—Canada, Mexico, the United States; South America—Brazil [64].

Lopharia ayresii (Berk. ex Cooke) Hjortstam, Mycotaxon 54: 188 (1995).

Specimens examined: on the trunk of angiosperm, 9 January 2019, CLZhao 10074, CLZhao 10,103 (SWFC).

Distribution: Africa—Angola, Eswatini, Kenya, Mauritius, South Africa, Tanzania, Uganda, Zimbabwe; Asia—China, Sri Lanka; North America—the United States; South America—Brazil, Ecuador; Oceania—Australia [64].

Lopharia mirabilis (Berk. and Broome) Pat., Bull. Soc. mycol. Fr. 11(1): 14 (1895).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3518, CLZhao 3519, CLZhao 3670; 6 October 2017, CLZhao 4399, CLZhao 4620; on the fallen branch of angiosperm, 4 January 2019, CLZhao 9376 (SWFC).

Distribution: Africa—Congo; Asia—China, India, Japan, Sri Lanka, Thailand; Europe—Russia [64].

Lyomyces bambusinus C.L. Zhao, in Chen and Zhao, MycoKeys 65: 105 (2020).

Specimens examined: on dead bamboo, 3 October 2017, CLZhao 3675; 11 January 2018, CLZhao 4808, CLZhao 4831, CLZhao 4840 (SWFC).

Distribution: Asia—China [64].

Lyomyces cremeus C.L. Zhao, in Chen and Zhao, MycoKeys 65: 108 (2020).

Specimens examined: on fallen branch of angiosperm, 21 August 2017, CLZhao 2812, CLZhao 4138; 23 August 2018, CLZhao 8295 (SWFC).

Distribution: Asia—China [64].

Lyomyces macrosporus C.L. Zhao, in Chen and Zhao, MycoKeys 65: 108 (2020).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4516, CLZhao 4531 (SWFC).

Distribution: Asia—China [64].

*Lyomyces wuliangshanensis* C.L. Zhao, in Chen and Zhao, MycoKeys 65: 111 (2020). Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4108, CLZhao 4144, CLZhao 4167; on fallen branch of angiosperm, 6 October 2017, CLZhao 4475;

on the stump of angiosperm, 5 October 2017, CLZhao 4206 (SWFC).

Distribution: Asia—China [64].

*Megasporia ellipsoidea* (B.K. Cui and P. Du) B.K. Cui and Hai J. Li, in Li and Cui, Mycologia 105(2): 375 (2013).

Specimens examined: on the fallen branch of angiosperm, 10 January 2019, CLZhao 10,360 (SWFC).

Distribution: Asia—China [64].

*Megasporia major* (G.Y. Zheng and Z.S. Bi) B.K. Cui and Hai J. Li, in Li and Cui, Mycologia 105(2): 375 (2013).

Specimens examined: on the fallen branch of angiosperm, 10 January 2019, CLZhao 10,363 (SWFC).

Distribution: Asia—China [64].

*Megasporoporiella pseudocavernulosa* B.K. Cui and Hai J. Li, in Li and Cui, Mycologia 105(2): 378 (2013).

Specimens examined: on the fallen branch of angiosperm, 4 January 2019, CLZhao 9339; 6 January 2019, CLZhao 9593, CLZhao 9688, CLZhao 9750; 7 January 2019, CLZhao 9806; 9 January 2019, CLZhao 10196, CLZhao 10,308 (SWFC).

Distribution: Asia—China [64].

*Megasporoporiella subcavernulosa* B.K. Cui and Hai J. Li, in Li and Cui, Mycologia 105(2): 378 (2013).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4442, CLZhao 4714 (SWFC).

Distribution: Asia—China [64].

*Metuloidea murashkinskyi* (Burt) Miettinen and Spirin, in Miettinen and Ryvarden, Ann. bot. fenn. 53(3-4): 165 (2016).

Specimens examined: on the trunk of angiosperm, 3 October 2017, CLZhao 3669, CLZhao 4454; on the fallen branch of angiosperm, 5 January 2019, CLZhao 9455 (SWFC).

Distribution: Asia—China, Japan; Europe—Bulgaria, Iran, Russia [64].

Microporus vernicipes (Berk.) Kuntze, Revis. gen. pl. (Leipzig) 3(3): 497 (1898).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4311, CLZhao 4361; 6 October 2017, CLZhao 4419; on the fallen branch of angiosperm, 2 October 2017, CLZhao 3672 (SWFC).

Distribution: Africa—Kenya, Zambia; Asia—China, Japan, Vietnam [64].

Microporus xanthopus (Fr.) Kuntze, Revis. gen. pl. (Leipzig) 3(3): 494 (1898).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4280; 5 January 2019, CLZhao 9577; on the fallen branch of angiosperm, 1 October 2017, CLZhao 3419 (SWFC).

Distribution: Asia—China, Japan, Malaysia [64].

*Neodatronia sinensis* B.K. Cui, Hai J. Li and Y.C. Dai, in Li, Cui and Dai, Persoonia 32: 178 (2014).

Specimens examined: on the fallen branch of angiosperm, 5 January 2019, CLZhao 9471 (SWFC).

Distribution: Asia—China [64].

Nigroporus vinosus (Berk.) Murrill, Bull. Torrey bot. Club 32(7): 361 (1905).

Specimens examined: on the trunk of angiosperm, 1 October 2017, CLZhao 3393; on the fallen branch of angiosperm, 1 October 2017, CLZhao 3376 (SWFC).

Distribution: Africa—Mozambique, Niger, Syria; Asia—China, Japan, Saudi Arabia; Europe—Spain; North America—Cuba, Mexico, Panama, the United States; Oceania—Australia [64].

*Oxychaete cervinogilva* (Jungh.) Miettinen, in Miettinen, Spirin, Vlasák, Rivoire, Stenroos and Hibbett, MycoKeys 17: 20 (2016).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3462, CLZhao 3465, CLZhao 3482, CLZhao 3642, CLZhao 3657, CLZhao 3663 (SWFC).

Distribution: Asia—China; Oceania—Australia [64].

Peniophora cinerea (Pers.) Cooke, Grevillea 8 (no. 45): 20 (1879).

Specimens examined: on the fallen branch of angiosperm, 3 January 2019, CLZhao 9254 (SWFC).

Distribution: Asia—China, India, Japan, Turkey; Europe—Austria, Denmark, Estonia, Finland, France, Germany, Italy, Netherlands, Norway, Poland, Portugal, Russia, Sweden, Ukraine; North America—Canada, Cuba, Jamaica, Mexico, the United States; South America—Brazil; Oceania—Australia, New Zealand [64].

Peniophora incarnata (Pers.) P. Karst., Hedwigia 28: 27 (1889) (Figure 7).



**Figure 7.** Basidiomata in situ of *Peniophora incarnata*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = (**D**) = 2 mm.

Specimens examined: on the trunk of angiosperm, 7 January 2019, CLZhao 9859 (SWFC).

Distribution: Africa—Morocco; Asia—China, India, Iran, Japan; Europe—Austria, Belgium, Czech Republic, Denmark, Finland, France, Germany, Iceland, Ireland, Italy, Netherlands, Norway, Poland, Portugal, Russia, Sweden, Switzerland, the United Kingdom, Ukraine; North America—Canada, Mexico, the United States; South America—Brazil, Chile; Oceania—Australia, New Zealand [64].

Peniophora pseudoversicolor Boidin, Bull. mens. Soc. linn. Lyon 34: 162 (1965).

Specimens examined: on the fallen branch of angiosperm, 4 January 2019, CLZhao 9415 (SWFC).

Distribution: Asia—China, Turkmenistan; Europe—Belgium, Denmark, Estonia, France, Germany, Italy, Portugal, Spain; Oceania—Australia [64].

Peniophorella fissurata C.L. Zhao, in Guan, Zhao and Zhao, Mycol. Progr. 19(4): 400 (2020).

Specimens examined: on the fallen branch of angiosperm, 6 October 2017, CLZhao 4539; 4 January 2019, CLZhao 9421 (SWFC).

Distribution: Asia—China [64].

Peniophorella pubera (Fr.) P. Karst., Bidr. Känn. Finl. Nat. Folk 48: 427 (1889).

Specimens examined: on the fallen branch of angiosperm, 4 January 2019, CLZhao 9336, CLZhao 9405 (SWFC).

Distribution: Asia—China, Japan; Europe—Denmark, Finland, Norway, Sweden, North America—Canada, the United States [64].

Peniophorella rude (Bres.) K.H. Larss., Mycol. Res. 111(2): 192 (2007).

Specimens examined: on the fallen branch of angiosperm, 1 October 2017, CLZhao 3360; 5 October 2017, CLZhao 4086, CLZhao 4300; 7 January 2019, CLZhao 9862; 9 January 2019, CLZhao 10212; 10 January 2019, CLZhao 10,419 (SWFC).

Distribution: Asia—China, Japan; North America—the United States; South America—Brazil [64].

*Perenniporia luteola* B.K. Cui and C.L. Zhao, in Zhao and Cui, Mycoscience 54(3): 235 (2013). Specimens examined: on the dead tree of angiosperm, 10 January 2019, CLZhao 10354; on the trunk of angiosperm, 10 January 2019, CLZhao 10334, CLZhao 10345, CLZhao 10403, CLZhao 10406, CLZhao 10411; on the fallen branch of angiosperm, 10 January 2019, CLZhao 10353, CLZhao 10355, CLZhao 10376, CLZhao 10384, CLZhao 10,393 (SWFC).

Distribution: Asia—China [64].

*Perenniporia mopanshanensis* C.L. Zhao, in Zhao and Ma, Mycotaxon 134(1): 132 (2019). Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4677,

CLZhao 4693; on the fallen branch of angiosperm, 6 October 2017, CLZhao 4708 (SWFC). Distribution: Asia—China [64].

*Phaeophlebiopsis peniophoroides* (Gilb. and Adask.) Floudas and Hibbett, Fungal Biology 119(7): 710 (2015).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3637 (SWFC).

Distribution: Asia—China; North America—the United States; South America— Brazil [64].

*Phanerochaete angustocystidiata* Sheng H. Wu, Bot. Bull. Acad. sin., Taipei 41(2): 166 (2000)

Specimens examined: on the fallen branch of angiosperm, 4 January 2019, CLZhao 9280, CLZhao 9505 (SWFC).

Distribution: Asia—China [64].

Phanerochaete australis Jülich, in Jermy, J. Linn. Soc., Bot. 81(1): 43 (1980).

Specimens examined: on the stump of angiosperm, 2 October 2017, CLZhao 3571 (SWFC). Distribution: Asia—China, Indonesia, Malaysia [64].

*Phanerochaete concrescens* Spirin and Volobuev, in Volobuev, Okun, Ordynets and Spirin, Mycol. Progr. 14(no. 80): 7 (2015).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4367; 5 January 2019, CLZhao 9585; 7 January 2019, CLZhao 9856, CLZhao 9899 (SWFC).

Distribution: Asia—China; Europe—Russia [64].

*Phanerochaete sanguineocarnosa* Floudas and Hibbett, Fungal Biology 119(7): 688 (2015). Specimens examined: on the trunk of angiosperm, 6 September 2017, CLZhao 4447, CLZhao 4639, CLZhao 4691 (SWFC).

Distribution: Asia—China; North America—the United States [64].

*Phanerochaete sordida* (P. Karst.) J. Erikss. and Ryvarden, in Eriksson, Hjortstam and Ryvarden, Cortic. N. Eur., 5 (Oslo): 1023 (1978).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4238, CLZhao 4270; 9 January 2019, CLZhao 10290; 10 January 2019, CLZhao 10347; on the fallen branch of angiosperm, 1 October 2017, CLZhao 3407; 5 October 2017, CLZhao 4082, CLZhao 4084, CLZhao 4091; 5 January 2019, CLZhao 9460; 7 January 2019, CLZhao 9936, CLZhao 9983; 9 January 2019, CLZhao 10172, CLZhao 10213, CLZhao 10314; 10 January 2019, CLZhao 10337, CLZhao 10366, CLZhao 10378, CLZhao 10390, CLZhao 10392, CLZhao

10402, CLZhao 10413, CLZhao 10439, CLZhao 10466, CLZhao 10518; on the fallen branch of *Pinus*, 5 October 2017, CLZhao 4087, CLZhao 4094, CLZhao 4134, CLZhao 4140, CLZhao 4187, CLZhao 4360 (SWFC).

Distribution: Asia—China, Iran, Japan, Nepal, Thailand; Europe—Denmark, Finland, Sweden; North America—Canada, the United States; Oceania—Australia [64].

*Phlebiopsis castanea* (Lloyd) Miettinen and Spirin, in Miettinen, Spirin, Vlasák, Rivoire, Stenroos and Hibbett, MycoKeys 17: 25 (2016).

Specimens examined: on the fallen branch of angiosperm, 2 October 2017, CLZhao 3501 (SWFC).

Distribution: Asia—China, Japan, South Korea, Thailand; Europe—Russia [64].

*Phlebiopsis crassa* (Lév.) Floudas and Hibbett, Fungal Biology 119(7): 710 (2015) (Figure 8).



**Figure 8.** Basidiomata in situ of *Phlebiopsis crassa*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = (**C**) = 1 cm; (**B**) = 1 mm; (**D**) = 2 mm.

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4128; 7 January 2019, CLZhao 9817; on the fallen branch of angiosperm, 1 October 2017, CLZhao 3339, 5 October 2017, CLZhao 4123, CLZhao 4172, CLZhao 4194, CLZhao 4225; 6 October 2017, CLZhao 4640; 4 January 2019, CLZhao 9316; 5 January 2019, CLZhao 9458, CLZhao 9539; 7 January 2019, CLZhao 9801, CLZhao 9827, CLZhao 9838 (SWFC).

Distribution: Asia—China, Vietnam; Europe—Belgium; North America—the United States; Oceania—Australia [64].

Pycnoporus sanguineus (L.) Murrill, Bull. Torrey bot. Club 31(8): 421 (1904).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4129, CLZhao 4234 (SWFC).

Distribution: Asia—China; Europe—France; North America—the United States; Oceania—Australia [64].

*Rhomboidia wuliangshanensis* C.L. Zhao, in Xu, Liu, Chen and Zhao, Mycotaxon 134(4): 656 (2020).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4406, CLZhao 4411 (SWFC).

Distribution: Asia—China [64].

*Rigidoporus pouzarii* Vampola and Vlasák, Czech Mycol. 64(1): 5 (2012).

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4524 (SWFC).

Distribution: Asia—China; Europe—Germany, Ukraine; North America—the United States [64].

*Serpula dendrocalami* C.L. Zhao, in Wang, Das, Bera, Chen, Bhatt, Ghosh, Hembrom, Hofstetter, Parihar, Vizzini, Xu, Zhao and Buyck, Cryptog. Mycol. 40(5): 88 (2019).

Specimens examined: on the dead bamboo, 2 October 2017, CLZhao 3521, CLZhao 3607, CLZhao 3626, CLZhao 3632 (SWFC).

Distribution: Asia—China [64].

*Sistotrema brinkmannii* (Bres.) J. Erikss., K. Fysiogr. Sällsk. Lund. Förhandl. 18 (no. 8): 134 (1948).

Specimens examined: on the trunk of angiosperm, 4 January 2019, CLZhao 9324, CLZhao 9344, CLZhao 9347 (SWFC).

Distribution: Asia—China; Europe—Bulgaria, Denmark, France, Germany, Italy, Poland, Portugal, Romania, Russia, Spain, the United Kingdom; North America—the United States; South America—Brazil [64].

*Sistotremastrum guttuliferum* Melo, M. Dueñas, Tellería and M.P. Martín, in Telleria, Melo, Dueñas, Salcedo, Beltrán-Tejera, Rodríguez-Armas and Martín, Mycol. Progr. 12(3): 688 (2012).

Specimens examined: on the trunk of angiosperm, 6 January 2019, CLZhao 9771; on the fallen branch of angiosperm, 5 January 2019, CLZhao 9454 (SWFC).

Distribution: Asia—China, South Korea; Europe—Romania, Spain [64].

Steccherinum bourdotii Saliba and A. David, Cryptog. Mycol. 9(2): 100 (1988).

Specimens examined: on the stump of angiosperm, 13 January 2018, CLZhao 4297; 10 January 2019, CLZhao 10,340 (SWFC).

Distribution: Asia—China, India; Europe—Belgium, Czech Republic, France, Germany, Russia, Switzerland [64].

Steccherinum confragosum Maas Geest. and Lang., Persoonia 8(2): 149 (1975).

Specimens examined: on the fallen branch of angiosperm, 5 October 2017, CLZhao 4347 (SWFC).

Distribution: Asia—China; Europe—Belarus, France [64].

Steccherinum ochraceum (Pers. ex J.F. Gmel.) Gray, Nat. Arr. Brit. Pl. (London) 1: 651 (1821). Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4523; 4 January 2019, CLZhao 9290; on the fallen branch of angiosperm, 7 January 2019, CLZhao 9953; 9 January 2019, CLZhao 10,055 (SWFC).

Distribution: Asia—China, Japan; Europe—Denmark, Germany, Norway, Spain, Sweden; North America—the United States; Oceania—Australia [64].

Terana coerulea (Lam.) Kuntze, Revis. gen. pl. (Leipzig) 2: 872 (1891).

Specimens examined: on the fallen branch of angiosperm, 6 January 2019, CLZhao 9613; 7 January 2019, CLZhao 9803, CLZhao 9812 (SWFC).

Distribution: Asia—China, Turkey; Europe—Belgium, Croatia, Czech Republic, Denmark, France, Germany, Greece, Italy, Macedonia, Portugal, Slovenia, Spain, Sweden, Switzerland, the United Kingdom, Ukraine [64].

*Tinctoporellus epimiltinus* (Berk. and Broome) Ryvarden, Trans. Br. mycol. Soc. 73(1): 18 (1979).

Specimens examined: on the trunk of angiosperm, 2 October 2017, CLZhao 3605 (SWFC).

Distribution: Asia—China, Japan, Malaysia; Europe—the United Kingdom; North America—Mexico [64].

Trametes ellipsospora Ryvarden, Mycotaxon 28(2): 539 (1987).

Specimens examined: on the fallen branch of angiosperm, 3 October 2017, CLZhao 3774 (SWFC).

Distribution: Asia—China; North America—the United States [64].

Trametes hirsuta (Wulfen) Lloyd, Mycol. Writ. (Cincinnati) 7(Letter 73): 1319 (1924)

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4278, CLZhao 4279, CLZhao 4342; on the fallen branch of angiosperm, 5 October 2017, CLZhao 4070, CLZhao 4170; 7 January 2019, CLZhao 9912, CLZhao 9986; on the stump of angiosperm, 2 October 2017, CLZhao 3572; on the fallen branch of *Pinus*, 5 October 2017, CLZhao 4295 (SWFC).

Distribution: Asia—China, India; Europe—Poland, the United Kingdom [64].

*Trametes pavonia* (Berk.) Fr., Nova Acta R. Soc. Scient. upsal., Ser. 3 1(1): 98 (1851). Specimens examined: on the fallen branch of angiosperm, 2 October 2017, CLZhao 3740 (SWFC).

Distribution: Asia—China; North America—the United States; South America—Argentina, Brazil [64].

Trametes versicolor (L.) Lloyd, Mycol. Notes (Cincinnati) 65: 1045 (1921) (Figure 9).



**Figure 9.** Basidiomata in situ of *Trametes versicolor*. (**A**,**B**) The front of the basidiomata; (**C**) the back of the basidiomata; (**D**) a section of hymenophore. Bars: (**A**) = 4 cm; (**B**) = (**D**) = 2 mm; (**C**) = 1 cm.

Specimens examined: on the trunk of angiosperm, 6 October 2017, CLZhao 4451, CLZhao 4497, CLZhao 4627; 5 January 2019, CLZhao 9587; 9 January 2019, CLZhao 10090; on the fallen branch of angiosperm, 6 October 2017, CLZhao 4487 (SWFC).

Distribution: Asia—China, India, Japan; Europe—Belgium, Denmark, Finland, France, Germany, Greece, Italy, Norway, Poland, Portugal, Spain, Sweden, the United Kingdom; North America—the United States [64].

Trametopsis cervina (Schwein.) Tomšovský, Czech Mycol. 60(1): 7 (2008).

Specimens examined: on the fallen branch of angiosperm, 5 January 2019, CLZhao 9449, CLZhao 9493 (SWFC).

Distribution: Africa—Morocco; Asia—China, India; Europe—Germany, the United Kingdom [64].

Trechispora nivea (Pers.) K.H. Larss., Symb. bot. upsal. 30(no. 3): 110 (1995).

Specimens examined: on the trunk of angiosperm, 9 January 2019, CLZhao 10154, CLZhao 10297; 10 January 2019, CLZhao 10373, CLZhao10389, CLZhao 10398, CLZhao 10,442 (SWFC).

Distribution: Asia—China; Europe—Austria, Belgium, Czech Republic, Denmark, Estonia, Finland, France, Germany, Italy, Norway, Poland, Romania, Russia, Spain, Sweden, Switzerland, the United Kingdom [64].

*Truncospora ochroleuca* (Berk.) Pilát, Sb. Nár. Mus. v Praze, Rada B, Prír. Vedy 9(2): 108 (1953).

Specimens examined: on the trunk of angiosperm, 5 October 2017, CLZhao 4350; on the stump of angiosperm, 4 January 2019, CLZhao 9328 (SWFC).

Distribution: Africa—Ethiopia, Kenya, Malawi, Tanzania; Asia—China, India, Japan, Malaysia, Singapore, Sri Lanka, Vietnam; Europe—Finland, France, Russia; North America—Mexico, the United States; South America—Brazil; Oceania—Australia, New Zealand [64].

## 4. Discussion

Both phyla Ascomycota and Basidiomycota cover around 97% of all fungal species in the kingdom fungi [65]. On the basis of the latest version of Ainsworth and Bisby's Dictionary of the Fungi [66], 1589 genera and more than 30 thousand species of Basidiomycota were included, and many taxa were recorded all over the world every year, which observably increase the fungal diversity [7,67–74]. The wood-decaying fungi have been extensively studied in China [6,7,68,72,73,75–78], but the diversity of wood-decaying fungi in the Wuliangshan area is yet elusive. The present paper is the first report of a series of studies devoted to wood-decaying fungi in this area and contributes to reveal abundance of wood-decaying species. A total of 2454 specimens belonging to 95 wood-decaying poroid and corticioid species were collected from the Wuliangshan area and identified based on morphological studies and phylogenetic analysis. According to the previous studies [6,16,63] combined with field observations, five species (5.26% of the total), Dacryobolus angiospermarum, Fibroporia albicans, Gloeophyllum sepiarium, Hymenochaete berteroi, and Serpula dendrocalami can cause a brown rot; the remaining 90 species (94.74% of the total) cause a white rot. In contrast, the proportion of brown-rot fungus in the Wuliangshan area is similar to that in the Haikou Forestry Farm which is 6.12% [78], but lower than Huangshan Mountains, Laojun Mountains and Great Xingan Mountains at 14.12%, 25% and 26.19%, respectively [75-77].

Although some remarkable explorations of wood-decaying fungi have been made from the Yunnan Province [57,68,79–89], only nine new species, Rhomboidia wuliangshanensis, Gloeodontia yunnanensis, Lyomyces bambusinus, L. cremeus, L. macrosporus, L. wuliangshanensis, Peniophorella fissurata, Perenniporia mopanshanensis and Serpula dendrocalami were found from the Wuliangshan area [85,86,90–93]. Species in the present list are mostly new to the studied area. All collected specimens were confirmed to be 95 species, which are distributed in 59 genera, 23 families, and 9 orders (Table 2). Of these 95 species listed, 67 species belong to the order Polyporales, accounting for 70.53%, indicating that polypores are the dominant group in this area; 12 species belong to the order Russulales; 6 species belong to the Hymenochaetales; 3 species belong to the Auriculariales; 2 species belong to Boletales and Trechisporales, and 1 species belongs to Cantharellales, Corticiales and Gloeophyllales. The present study indicates that Aleurodiscus botryosus, A. cerussatus, A. mirabilis, Asterostroma muscicola, Auricularia cornea, A. delicata, Bjerkandera adusta, Byssomerulius corium, Daedaleopsis confragosa, Datronia stereoides, Fomes fomentarius, Hydnoporia corrugata, Lenzites betulinus, Lopharia ayresii, Nigroporus vinosus, Peniophora cinerea, P. incarnata, Sistotrema brinkmannii, Steccherinum ochraceum, Terana coerulea, Trametes versicolor, Trechispora nivea and Truncospora ochroleuca species are common, Abundisporus quercicola, Aleurodiscus isabellinus, Gloeodontia yunnanensis, Lyomyces bambusinus, L. cremeus, L. macrosporus, L. wuliangshanensis, Megasporoporiella pseudocavernulosa, Peniophorella fissurata, Perenniporia luteola, P. mopanshanensis, Phanerochaete angustocystidiata, Rhomboidia wuliangshanensis and Serpula dendrocalami are rare worldwide according to the distribution, whereas 69 species belong to rare native species (fewer than five specimens followed previous study [6]); 49 species are distributed in Eurasia, and 61 are spread in the Northern Hemisphere; Bjerkandera adusta and Fomes fomen*tarius* could function as indicators that detect heavy metal biosorption; *Bjerkandera adusta*, Daedaleopsis confragosa, Datronia stereoides, Fomes fomentarius and Peniophora incarnata are expansive, and Cerrena albocinnamomea, Heterobasidion insulare, Phanerochaete concrescens and Phlebiopsis castanea are invasive in the Wuliangshan area. In addition, the underlying fungal diversity in the Wuliangshan area is unpredictable, and a more intensive investigation is needed to complement the further biodiversity.

Ordore Familio Conora

Table 2. Number of wood-inhabiting poroid and corticioid species in the most representative orders, families, and genera in the study area and proportion accounting for total species number.

As an essential component of the forest ecosystems, most of the wood-inhabiting poroid and corticioid fungi function as wood decomposers that release matter and energy to the ecological system [94–96]. Herein, we further found 2 edible fungi, Auricularia cornea and A. delicata, according to Dai et al. [97], together with 14 medicinal fungi, Bjerkandera adusta, Cerrena zonata, Daedaleopsis confragosa, Earliella scabrosa, Fomes fomentarius, Ganoderma australe, Gloeophyllum sepiarium, Hexagonia tenuis, Irpex lacteus, Lenzites betulinus, Lopharia ayresii, *Pycnoporus sanguineus, Trametes hirsuta* and *T. versicolor*, according to Dai and Yang [98]. Some notable medical fungi were found, such as *Bjerkandera adusta*, a potential antineoplastic agent [99], as well as Trametes hirsute, which possesses the effect of curing rheumatism and relieving cough [100,101]. In addition, an industrial white-rot fungus was also found in this study. As a target of interest, Trametes versicolor was conducted under a free or immobilized laccase condition exploring its biocatalytic characterization in the activation zone [102]. Moreover, though these saprophytic species possess powerful enzymes, which can effectively degrade lignocellulose [103], Abundisporus quercicola, Bjerkandera adusta, Ceriporia lacerata, Cerrena zonata, C. albocinnamomea, Daedaleopsis confragosa, Fomes fomentarius and Gyrodontium sacchari are treated as forest pathogens contrarily [104].

Phylogenetically, based on a study of the molecular systematics on the subphyla Agaricomycotina, Pucciniomycotina and Ustilaginomycotina with combined nLSU, SSU, 5.8s, rpb1, rpb2, and tef1 datasets, He et al. [105] revealed 1928 currently used genera names, which are distributed in 241 families, 68 orders, 18 classes and 4 subphyla. In the present study, 95 species nested in 59 genera, 23 families, and 9 orders based on the ITS dataset, which is similar to the previous studied topology [105], were revealed. Sequences of ITS gene regions of the studied samples were generated, which illustrates that all collected fungal specimens from the Wuliangshan area belonging to Agaricomycetes distribute into Auriculariales, Boletales, Cantharellales, Corticiales, Gloeophyllales, Hymenochaetales, Polyporales, Russulales and Trechisporales (Figure 1). Phylogenetic analyses of all studied samples shows that the individual taxon clusters closely with the downloaded reliable sequence from previous studies with a supported rate (Figure 2).

Based on the early embrace of molecular systematics by mycologists, both the discovery and classification of fungi among the more basal branches of the tree are now coming to light from genomic analyses and environmental DNA surveys [1]. Several

Oldels					1 annines				Genera		
Order	Family	Genera	Spp.	%	Family	Genera	Spp.	%	Genera	Spp.	%
Polyporales	9	40	67	70.53	Polyporaceae	18	28	29.47	Aleurodiscus	5	5.26
Russulales	5	6	12	12.63	Phanerochaetaceae	6	11	11.58	Phanerochaete	5	5.26
Hymenochaetales	2	3	6	6.32	Steccherinaceae	5	8	8.42	Hyphoderma,	4	4.21
Auriculariales	1	1	3	3.16	Hyphodermataceae	e 2	8	8.42	Lyomyces	4	4.21
Boletales	2	4	2	2.11	Stereaceae	2	6	6.32	Trametes	4	4.21
Trechisporales	1	2	2	2.11	Irpicaceae	5	5	5.26	Auricularia	3	3.16
Cantharellales	1	1	1	1.05	Cerrenaceae	2	3	3.16	Peniophora	3	3.16
Corticiales	1	1	1	1.05	Hymenochaetaceae	e 2	3	3.16	Peniophorella	3	3.16
Gloeophyllales	1	1	1	1.05	Rickenellaceae	1	3	3.16	Steccherinum	3	3.16
1 5					Subtotal	43	75	78.95	Subtotal	34	35.79
					Other families				Other		
				(14)	16	20	21.05	genera (50)	61	64.21	
Total	23	59	95	100%	23	59	95	100%	59	95	100%

comprehensively remarkable collections of wood-decaying fungi in China have been performed based on the combination of morphological features and molecular evidence; Dai illustrated 80 pathogenic wood-decaying fungi in 2005, which mainly belong to Ganodermataceae, Climacodontaceae, Corticiaceae, Hericiaceae, Hymenochataceae, Polyporaceae, and Schizophyllaceae of Basidiomycota [106]. In addition, Dai reported 140 wood-decaying fungi on stored wood or structural timber in 2009, and 29 are brown-rot and 111 are white-rot fungus [107]. The fungal diversity from the Hainan area in China showed that it includes 240 species, of which 199 are polypores, and 41 are corticoid and other woodinhabiting fungi [108]. In addition, this place is located at the longitude and latitude of 108°37′–111°03′ E and 18°10′–20°10′ N. The Atlas of Chinese Macrofungal Resources revealed that it has 1819 species belonging to 509 genera. According to their morphological characteristics, they are divided into 10 groups, including 196 larger Ascomycetes, 21 jelly fungi, 47 coral fungi, 637 polyporoid, hydnaceous and thelephoroid fungi, 11 cantharelloid fungi, 653 agarics, 130 boletes, 75 gasteroid fungi, 16 larger pathogenic fungi on crops, and 33 larger Myxomycetes [109]. The book of Atlas of Macrofungi in Saihanwula National Nature Reserve of Inner Mongolia indicated that it includes 310 species, located in 3 phyla, 8 classes, 24 orders, 66 families, 152 genera [110]; its longitude and latitude are 118°18′–118°55′ E and 43°59′–44°27′ N. The result shows that the tropical and subtropical areas have enriched fungal diversity, which supports our research regarding to the richness of wood-decaying fungi in China.

In the last 20 years, dramatic changes in higher-level taxonomy have occurred, as evidenced by a tripling of fungal phyla from 4 to 12. The hidden and microscopic nature of many fungi also means that their biodiversity is undersampled, in which perhaps less than 5% of the estimated two to four million species have been formally described [2,16,111]. The present work will comprehensively improve the understanding of the diversity of wood-decaying fungi in this area, in which 29 species are new for China, i.e., Aleurodiscus botryosus, Aleurodiscus isabellinus, Auricularia asiatica, Crustodontia chrysocreas, Flavodon ambrosius, Gloeodontia yunnanensis, Hyphoderma moniliforme, Hyphoderma nudicephalum, Hyphoderma transiens, Laxitextum incrustatum, Lyomyces bambusinus, Lyomyces cremeus, Lyomyces macrosporus, Lyomyces wuliangshanensis, Peniophora pseudoversicolor, Peniophorella fissurata, Peniophorella pubera, Peniophorella rude, Perenniporia mopanshanensis, Phaeophlebiopsis peniophoroides, Phanerochaete concrescens, Phanerochaete sanguineocarnosa, Rhomboidia wuliangshanensis, Rigidoporus pouzarii, Serpula dendrocalami, Sistotremastrum guttuliferum, Steccherinum bourdotii, Steccherinum confragosum and Terana coerulea. The research region of woodinhabiting fungi in the Wuliangshan area covers 2,581 square kilometres, accounting for 0.65% of the area of the Yunnan Province and for 0.027% of China. This work is conducive to the rational utilization and effective protection of fungal resources, and provides scientific basis for the prevention and control of forest diseases in this area.

Author Contributions: Conceptualization, C.-L.Z.; methodology, C.-L.Z. and X.H.; software, C.-L.Z. and X.H.; validation, C.-L.Z. and X.H.; formal analysis, C.-L.Z. and X.H.; investigation, C.-L.Z. and X.H.; resources, C.-L.Z.; writing—original draft preparation, C.-L.Z. and X.H.; writing—review and editing, C.-L.Z. and X.H.; visualization, C.-L.Z. and X.H.; supervision, C.-L.Z.; project administration, C.-L.Z.; funding acquisition, C.-L.Z. All authors have read and agreed to the published version of the manuscript.

**Funding:** The research was supported by the National Natural Science Foundation of China (Project No. 32170004, U2102220), Yunnan Fundamental Research Project (Grant No. 202001AS070043) and High-level Talents Program of Yunnan Province (YNQR-QNRC-2018-111).

Institutional Review Board Statement: Not applicable for studies involving humans or animals.

Informed Consent Statement: Not applicable for studies involving humans.

**Data Availability Statement:** Publicly available datasets were analyzed in this study. This data can be found here: https://www.ncbi.nlm.nih.gov/; https://www.mycobank.org/page/Simple%20nam es%20search; http://purl.org/phylo/treebase, submission ID 29040; accessed on 23 November 2021.

Conflicts of Interest: The authors declare no conflict of interest.

## References

- 1. James, T.Y.; Stajich, J.E.; Hittinger, C.T.; Rokas, A. Toward a fully resolved fungal tree of life. *Annu. Rev. Microbiol.* **2020**, *74*, 291–313. [CrossRef] [PubMed]
- 2. Blackwell, M. The Fungi: 1, 2, 3 ... 5.1 million species? Am. J. Bot. 2011, 98, 426–438. [CrossRef] [PubMed]
- 3. Taylor, D.L.; Hollingsworth, T.N.; Mcfarland, J.W. A first comprehensive census of fung i in soil reveals bothhyperdiversity and fine-scale niche partitioning. *Ecol. Monogr.* **2014**, *84*, 3–20. [CrossRef]
- 4. Hibbett, D. The invisible dimension of fungal diversity. *Science* 2016, 351, 1150. [CrossRef]
- 5. Russell, R.; Paterson, M. Ganoderma—A therapeutic fungal biofactory. Phytochemistry 2006, 67, 1985–2001. [CrossRef]
- 6. Dai, Y.C. Polypore diversity in China with an annotated checklist of Chinese polypores. Mycoscience 2012, 53, 49–80. [CrossRef]
- 7. Dai, Y.C.; Cui, B.K.; Si, J.; He, S.H.; Hyde, K.D.; Yuan, H.S.; Liu, X.Y.; Zhou, L.W. Dynamics of the worldwide number of fungi with emphasis on fungal diversity in China. *Mycol. Prog.* **2015**, *14*, 62. [CrossRef]
- 8. Vinay, D.S.; Ryan, E.P.; Pawelec, G.; Talib, W.H.; Stagg, J.; Elkord, E.; Lichtor, T.; Decker, W.K.; Whelan, R.L.; Kumara, S.; et al. Immune evasion in cancer: Mechanistic basis and therapeutic strategies. *Semin. Cancer Biol.* **2015**, *35*, S185–S198. [CrossRef]
- 9. M'Barek, H.N.; Arif, S.; Taidi, B.; Hajjaj, H. Consolidated bioethanol production from olive mill waste: Wood-decay fungi from central Morocco as promising decomposition and fermentation biocatalysts. *Biotechnol. Rep.* 2020, *28*, e00541. [CrossRef]
- 10. Runnel, K.; Miettinen, O.; Lohmus, A. Polypore fungi as a flagship group to indicate changes in biodiversity—A test case from Estonia. *IMA Fungus* **2021**, *12*, 2. [CrossRef]
- 11. Yu, Q.G. Wuliangshan National Nature Reserve; Yunnan Science and Technology Press: Kunming, China, 2004; pp. 103–105, 118, 197.
- 12. Lu, S.Z.; Yu, Q.G.; Cao, S.W. Measurement of ecosystem diversity in Wuliangshan National Nature Reserve and its adjacent area in Yunnan Province. *J. Anhui Agric. Sci.* 2008, *36*, 2426–2428, 2435.
- 13. Luo, Z.H.; Xie, Y.N.; Luo, Y.; Li, X.Y.; Lu, C.R. Population Characteristics of *Tsuga dumosa* in Wuliangshan National Nature Reserve. *J. West China For. Sci.* **2013**, *42*, 94–99.
- 14. Gao, L.L.; Yang, H.J.; Liu, G.Z.; Wang, C.H.; Yang, G.P.; Cai, C.T. Diversity of medicinal vascular plant resources in Wuliangshan National Nature Reserve, Yunnan, China. *Plant Sci. J.* **2018**, *36*, 203–210.
- 15. Peng, H.; Wu, Z.Y. The Preliminary Floristical study on mid-montane humid evergreen broad-leaved forest in Mt. Wuliangshan. *Acta Bot. Yunnanica* **1998**, *20*, 12–22.
- 16. Dai, Y.C.; Yang, Z.L.; Cui, B.K.; Wu, G.; Yuan, H.S.; Zhou, L.W.; He, S.H.; Ge, Z.W.; Wu, F.; Wei, Y.L.; et al. Diversity and systematics of the important macrofungi in Chinese forests. *Mycosystema* **2021**, *40*, 770–805. [CrossRef]
- 17. Petersen, J.H. *Farvekort. The Danish Mycological Society's Colour-Chart;* Foreningen til Svampekundskabens Fremme: Paris, France, 1996; pp. 1–6.
- 18. White, T.J.; Bruns, T.; Lee, S.; Taylor, J. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protoc. Guide Methods Appl.* **1990**, *18*, 315–322. [CrossRef]
- 19. Xu, T.M. Study on the Resources and Taxonomy of Wood-Decaying Fungi in Wuliangshan National Nature Reserve, Yunnan Province; Southwest Forestry University: Kunming, China, 2020.
- 20. Vu, D.; Groenewald, M.; De Vries, M.; Gehrmann, T.; Stielow, B.; Eberhardt, U.; Al-Hatmi, A.; Groenewald, J.Z.; Cardinali, G.; Houbraken, J.; et al. Large-scale generation and analysis of filamentous fungal DNA barcodes boosts coverage for kingdom fungi and reveals thresholds for fungal species and higher taxon delimitation. *Stud. Mycol.* **2019**, *92*, 135–154. [CrossRef]
- 21. Tian, Y.; Ghobad-Nejhad, M.; He, S.H.; Dai, Y.C. Three new species of *Aleurodiscus* s.l. (Russulales, Basidiomycota) from southern China. *MycoKeys* **2018**, *37*, 93–107. [CrossRef]
- 22. Asl, A.; Syo, A.; Msp, A.; Kim, T.; Yoo, J.S.; Seelan, J.S.S.; Lim, Y.W. Co-occurrence patterns of wood-decaying fungi and ants in dead pines of South Korea—ScienceDirect. J. Asia Pac. Entomol. 2019, 22, 1154–1160. [CrossRef]
- 23. Bandara, A.R.; Karunarathna, S.C.; Phillips, A.J.L.; Mortimer, P.E.; Xu, J.C.; Kakumyan, P.; Hyde, K.D. Diversity of *Auricularia* (Auriculariaceae, Auriculariales) in Thailand. *Phytotaxa* 2017, 292, 19. [CrossRef]
- 24. Wu, F.; Yuan, Y.; Rivoire, B.; Dai, Y.C. Phylogeny and diversity of the *Auricularia mesenterica* (Auriculariales, Basidiomycota) complex. *Mycol. Prog.* **2015**, *14*, 42. [CrossRef]
- 25. Floudas, D.; Hibbett, D.S. Revisiting the taxonomy of *Phanerochaete* (Polyporales, Basidiomycota) using a four gene dataset and extensive ITS sampling. *Fungal Biol.* **2015**, *119*, 679–719. [CrossRef]
- Águeda, B.; Parladé, J.; Fernandez-Toiran, L.M.; Cisneros, Ó.; Miguel, A.M.; Modrego, M.P.; Martínez-Peña, F.; Pera, J. Mycorrhizal synthesis between *Boletus edulis* species complex and rockroses (*Cistus* sp.). *Mycorrhiza* 2008, 18, 443–449. [CrossRef]
- 27. Deo, G.S.; Khatra, J.; Buttar, S.; Wai, M.L.; Chow, H.L. Anti-proliferative, immuno-stimulatory and anti-inflammatory activities of extracts derived from mushrooms collected in Haida Gwaii, BC (Canada). *Int. J. Med. Mushrooms* 2019, 21, 629–643. [CrossRef]
- Shirouzu, T.; Hosaka, K.; Nam, K.O.; Weir, B.S.; Hosoya, T. Phylogenetic relationships of eight new Dacrymycetes collected from New Zealand. *Persoonia* 2017, 38, 156–169. [CrossRef]

- Shirouzu, T.; Hirose, D.; Oberwinkler, F.; Shimomura, N.; Maekawa, N.; Tokumasu, S. Combined molecular and morphological data for improving phylogenetic hypothesis in Dacrymycetes. *Mycologia* 2013, 105, 1110–1125. [CrossRef]
- 30. Zamora, J.C.; Ekman, S. Phylogeny and character evolution in the Dacrymycetes, and systematics of Unilacrymaceae and Dacryonaemataceae fam. nov. *Persoonia* 2020, 44, 161–205. [CrossRef]
- Chen, Y.Y.; Li, H.J.; Cui, B.K. Molecular phylogeny and taxonomy of *Fibroporia* (Basidiomycota) in China. *Phytotaxa* 2015, 203, 47–54. [CrossRef]
- 32. Li, Y.; Bateman, C.C.; Skelton, J.; Jusino, M.A.; Nolen, Z.J.; Simmons, D.R.; Hulcr, J. Wood decay fungus *Flavodon ambrosius* (Basidiomycota: Polyporales) is widely farmed by two genera of ambrosia beetles. *Fungal Biol.* **2017**, 121, 984–989. [CrossRef]
- 33. Jeppson, M.; Nilsson, R.H.; Larsson, E. European earthstars in Geastraceae (Geastrales, Phallomycetidae)—A systematic approach using morphology and molecular sequence data. *Syst. Biodivers.* **2013**, *11*, 437–465. [CrossRef]
- 34. Larsson, K.H. Re-thinking the classification of corticioid fungi. Mycol. Res. 2007, 111, 1040–1063. [CrossRef] [PubMed]
- Yuan, Y.; Chen, J.J.; Korhonen, K.; Martin, F.; Dai, Y.C. An updated global species diversity and phylogeny in the forest pathogenic genus *Heterobasidion* (Basidiomycota, Russulales). *Front. Microbiol.* 2021, *11*, 596393. [CrossRef] [PubMed]
- Ji, X.; Wu, D.M.; Song, C.G.; Liu, S.; Si, J.; Cui, B.K. Two new *Neofomitella* species (Polyporaceae, Basidiomycota) based on morphological and molecular evidence. *Mycol. Prog.* 2019, 18, 593–602. [CrossRef]
- Li, H.J.; Cui, B.K.; Dai, Y.C. Taxonomy and multi-gene phylogeny of *Datronia* (Polyporales, Basidiomycota). *Persoonia* 2014, 32, 170–182. [CrossRef]
- 38. Yurchenko, E.; Wu, S.H. *Hyphoderma moniliforme* and *H. nemorale* (Basidiomycota) newly recorded from China. *Mycosphere* **2015**, *6*, 113–121. [CrossRef]
- 39. Larsson, K.H. Molecular phylogeny of *Hyphoderma* and the reinstatement of *Peniophorella*. *Mycol. Res.* **2007**, *111*, 186–195. [CrossRef]
- Sulzbacher, M.A.; Grebenc, T.; Nouhra, E.R.; Giachini, A.J.; Martín, M.P.; Baseia, L.G. *Hysterangium atlanticum* sp. nov. forms ectomycorrhizae with Coccoloba species (Polygonaceae) from the Atlantic rainforest of Northeastern Brazil. *Symbiosis* 2019, 78, 275–286. [CrossRef]
- Miettinen, O.; Larsson, E.; Sjokvist, E.; Larsson, K.H. Comprehensive taxon sampling reveals unaccounted diversity and morphological plasticity in a group of dimitic polypores (Polyporales, Basidiomycota). *Cladistics Int. J. Willi Hennig Soc.* 2012, 28, 251–270. [CrossRef]
- Mudalungu, C.M.; Richter, C.; Wittstein, K.; Abdalla, M.A.; Matasyoh, J.C.; Stadler, M.; Süssmuth, R.D. Laxitextines A and B, Cyathane Xylosides from the Tropical Fungus *Laxitextum incrustatum*. J. Nat. Prod. 2016, 79, 894–898. [CrossRef]
- 43. Liu, S.L.; Nakasone, K.K.; Wu, S.H.; He, S.H.; Dai, Y.C. Taxonomy and phylogeny of *Lopharia* s.s *Dendrodontia*, *Dentocorticium* and *Fuscocerrena* (Basidiomycota, Polyporales). *MycoKeys* **2018**, *32*, 25–48. [CrossRef]
- Miettinen, O.; Spirin, V.; Josef, V.; Rivoire, B.; Hibbett, D. Polypores and genus concepts in Phanerochaetaceae (Polyporales, Basidiomycota). *MycoKeys* 2016, 17, 1–46. [CrossRef]
- Hallenberg, N.; Nilsson, R.H.; Antonelli, A.; Wu, S.H.; Maekawa, N.; Nordén, B. The *Peniophorella praetermissa* species complex (Basidiomycota). *Mycol. Res.* 2008, 111, 1366–1376. [CrossRef]
- 46. Cabral, T.S.; Silva, B.D.; María, P.M.; Clement, C.R.; Hosaka, K.; Baseia, I.G. Behind the veil—Exploring the diversity in *Phallus indusiatus* s.l. (Phallomycetidae, Basidiomycota). *MycoKeys* **2019**, *58*, 103–127. [CrossRef]
- 47. Volobuev, S.; Okun, M.; Ordynets, A.; Spirin, V. The *Phanerochaete sordida* group (Polyporales, Basidiomycota) in temperate Eurasia, with a note on *Phanerochaete pallida*. *Mycol. Prog.* **2015**, *14*, 80. [CrossRef]
- Melera, S.; Ostellari, C.; Roemer, N.; Avis, P.G.; Tonolla, M.; Barja, F.; Narduzzi-Wicht, B. Analysis of morphological, ecological and molecular characters of *Russula pectinatoides* Peck and *Russula praetervisa* Sarnari, with a description of the new taxon *Russula* recondita Melera & Ostellari. Mycol. Prog. 2016, 16, 117–134. [CrossRef]
- 49. Palmer, J.M.; Lindner, D.L.; Volk, T.J. Ectomycorrhizal characterization of an American chestnut (Castanea dentata)-dominated community in Western Wisconsin. *Mycorrhiza* **2008**, *19*, 27–36. [CrossRef]
- 50. Alshammari, N.; Stephenson, S.L. A preliminary study of wood-decay fungi in forests of northwest Arkansas. *Curr. Res. Environ. Appl. Mycol.* **2018**, *8*, 556–563. [CrossRef]
- Justo, A.; Miettinen, O.; Floudas, D.; Ortiz-Santana, B.; Sjökvist, E.; Lindner, D.; Nakasone, K.; Niemelä, T.; Larsson, K.H.; Ryvarden, L.; et al. A revised family-level classification of the Polyporales (Basidiomycota). *Fungal Biol.* 2017, 121, 798–824. [CrossRef]
- 52. Zhao, C.L.; Cui, B.K.; Steffen, K.T. Yuchengia, a new polypore genus segregated from *Perenniporia* (Polyporales) based on morphological and molecular evidence. *Nord. J. Bot.* 2013, *31*, 331–338. [CrossRef]
- 53. Milenkovic, I.; Tomsovsky, M.; Karadzic, D.; Veselinovic, M. Decline of *Paulownia tomentosa* caused by *Trametes hirsuta* in Serbia. *For. Pathol.* **2018**, *48*, e12438. [CrossRef]
- Justo, A.; Hibbett, D.S. Phylogenetic classification of *Trametes* (Basidiomycota, Polyporales) based on a five-marker dataset. *Taxon* 2011, 60, 1567–1583. [CrossRef]
- Telleria, M.T.; Melo, I.; Duenas, M.; Larsson, K.H.; Martín, M.P.P. Molecular analyses confirm *Brevicellicium* in Trechisporales. *IMA Fungus* 2013, 4, 21–28. [CrossRef]
- Hall, T.A. Bioedit: A user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. Nucleic Acids Symp. Ser. 1999, 41, 95–98. [CrossRef]

- 57. Zhao, C.L.; Wu, Z.Q. *Ceriporiopsis kunmingensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analysis. *Mycol. Prog.* 2017, *16*, 93–100. [CrossRef]
- 58. Swofford, D.L. *PAUP\*: Phylogenetic Analysis Using Parsimony (\*and Other Methods);* Version 4.0b10; Sinauer Associates: Sunderland, MA, USA, 2002.
- 59. Felsenstein, J. Confidence intervals on phylogenetics: An approach using bootstrap. Evolution 1985, 39, 783–791. [CrossRef]
- 60. Miller, M.A.; Pfeiffer, W.; Schwartz, T. The CIPRES Science Gateway: Enabling High-Impact Science for Phylogenetics Researchers with Limited Resources. *Assoc. Comput. Mach.* 2012, 39, 1–8. [CrossRef]
- 61. Nylander, J.A.A. *MrModeltest v2. Program Distributed by the Author*; Evolutionary Biology Centre, Uppsala University: Uppsala, Sweden, 2004.
- 62. Ronquist, F.; Huelsenbeck, J.P. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **2003**, *19*, 1572–1574. [CrossRef]
- 63. Dai, Y.C. A revised checklist of corticioid and hydnoid fungi in China for 2010. Mycoscience 2011, 52, 69–79. [CrossRef]
- 64. Zhao, C.L. Wood Inhabiting Fungi Diversity in Wuliangshan Area; China Forestry Publishing House: Beijing, China, 2021.
- 65. Willis, K.J. State of the World's Fungi 2018; Report; Royal Botanic Gardens: Kew, UK, 2018.
- 66. Kirk, P.M.; Cannon, P.F.; Minter, D.W.; Stalpers, J.A. *Ainsworth & Bisby's Dictionary of the Fungi*, 10th ed.; CAB International: Wallingford, UK, 2008.
- Ayesha, P.; Maham, A.; Ariza, A.; Husnain, H. "Ganoderma lucidum sensu lato"—A sacred mushroom for immortality. Stud. Fungi 2020, 5, 508–516. [CrossRef]
- 68. Huang, R.X.; Luo, K.Y.; Zhao, C.L. *Phlebia nigrodontea* sp. nov. in Meruliaceae (Polyporales) with a black hymenial surface. *Phytotaxa* **2020**, 458, 195–206. [CrossRef]
- 69. Himani, S.; Krishnappa, M. *Xylaria oxyacanthae* (Xylariaceae), a new record on Diospyros melanoxylon from India. *Stud. Fungi* **2020**, *5*, 485–490. [CrossRef]
- 70. Blanco-Dios, J.B. Notes on the genus *Entoloma* s.l. in the Northwest of the Iberian Peninsula (XI): A new species in subgenus *Entoloma*. *Stud. Fungi* **2021**, *6*, 64–70. [CrossRef]
- Kumar, S.; Singh, R. Global diversity and distribution of distoseptosporic micromycete *Corynespora* Güssow (Corynesporascaceae): An updated checklist with current status. *Stud. Fungi* 2021, *6*, 1–63. [CrossRef]
- 72. Wang, H.; Zhao, C.L. *Hyphodermella aurantiaca* sp. nova. (Polyporales, Basidiomycota) as evidenced by morphological characters and phylogenetic analyses. *Ann. Bot. Fenn.* **2021**, *58*, 61–68. [CrossRef]
- Zong, T.K.; Liu, C.M.; Wu, J.R.; Zhao, C.L. Trechispora daweishanensis and T. xantha spp. nov. (Hydnodontaceae, Trechisporales) found in Yunnan Province of China. Phytotaxa 2021, 479, 147–159. [CrossRef]
- 74. Boonyuen, N.; Chuaseeharonnachai, C.; Nuankaew, S.; Kwantong, P.; Pornputtapong, N.; Suwannarach, N.; Jones, E.B.G.; Somrithipol, S. Novelties in Fuscosporellaceae (Fuscosporellales): Two New *Parafuscosporella* from Thailand Revealed by Morphology and Phylogenetic Analyses. *Diversity* **2021**, *13*, 517. [CrossRef]
- Cui, B.K.; Jia, B.S. Wood-rotting fungi in eastern China 7. Polypores from Huangshan Mountain, Anhui Province. *Mycosystema* 2011, 30, 529–535. [CrossRef]
- 76. Cui, B.K.; Yu, C.J. Fungal flora and population structure of polypores in the Great Hingan Mountains. *Acta Ecol. Sin.* **2011**, *31*, 3700–3709.
- Cui, B.K.; Dai, Y.C. Wood-decaying fungi in eastern Himalayas 3. Polypores from Laojunshan Mountains, Yunnan Province. *Mycosystema* 2012, 31, 486–492.
- 78. He, X.; Chen, J.Z.; Zhao, C.L. Diversity of wood-decaying fungi in Haikou Forestry Farm, Yunnan Province, P.R. China. *Stud. Fungi* **2021**, *6*, 365–377. [CrossRef]
- 79. Yuan, H.S.; Dai, Y.C. Polypores from northern and central Yunnan Province, Southwestern China. Sydowia 2008, 60, 147–159.
- Wu, Z.Q.; Shen, S.; Luo, K.Y.; Wang, Z.H.; Zhao, C.L. Morphological and molecular identification of a new species of *Atraporiella* (Polyporales, Basidiomycota) in China. *Phytotaxa* 2017, 332, 31–40. [CrossRef]
- Shen, S.; Ma, X.; Xu, T.M.; Zhao, C.L. *Phlebia ailaoshanensis* sp. nov. (Polyporales, Basidiomycota) evidenced by morphological characters and phylogenetic analyses. *Phytotaxa* 2018, 373, 184–196. [CrossRef]
- 82. Wu, Z.Q.; Xu, T.M.; Shen, S.; Liu, X.F.; Luo, K.Y.; Zhoa, C.L. *Elaphroporia ailaoshanensis* gen. et sp nov in Polyporales (Basidiomycota). *MycoKeys* **2018**, *29*, 81–95. [CrossRef]
- 83. Liu, X.F.; Shen, S.; Zhao, C.L. Morphological and molecular identification of a new species of *Eichleriella* (Auriculariales, Basidiomycota) in China. *Phytotaxa* **2019**, 404, 245–254. [CrossRef]
- Luo, K.Y.; Ma, X.; Zhao, C.L. Neofavolus yunnanensis sp. nov. (Polyporales, Basidiomycota) from China: Evidence from morphology and DNA sequence data. Phytotaxa 2019, 408, 109–116. [CrossRef]
- Xu, T.M.; Liu, X.F.; Chen, Y.H.; Zhao, C.L. *Rhomboidia wuliangshanensis* gen. et sp. nov. from southwestern China. *Mycotaxon* 2019, 134, 649–662. [CrossRef]
- 86. Chen, J.Z.; Zhao, C.L. Morphological and molecular identification of four new resupinate species of *Lyomyces* (Hymenochaetales) from southern China. *MycoKeys* **2020**, *65*, 101–118. [CrossRef]
- 87. Wang, R.Z.; Yuan, Z.Y.; Han, C.Z.; Zhao, C.L. Morphological and molecular identification of a new species of *Cinereomyces* (Polyporales, Basidiomycota) in southern China. *Phytotaxa* **2020**, 459, 51–60. [CrossRef]
- 88. Gu, Z.R.; Zhao, C.L. The hidden wood-decaying fungal diversity: Rhizochaete from east Asia. Diversity 2021, 3, 503. [CrossRef]

- 89. Luo, K.Y.; Qu, M.H.; Zhao, C.L. Additions to the knowledge of corticioid *Xylodon* (Schizoporaceae, Hymenochaetales): Introducing three new *Xylodon* species from southern China. *Diversity* **2021**, *13*, 581. [CrossRef]
- Wang, X.H.; Das, K.; Bera, I.; Chen, Y.H.; Bhatt, R.P.; Ghosh, A.; Hembrom, M.E.; Hofstetter, V.; Parihar, A.; Vizzini, A.; et al. Fungal Biodiversity Profiles 81–90. *Cryptogam. Mycol.* 2019, 40, 57–95. [CrossRef]
- 91. Zhao, C.L.; Ma, X. Perenniporia mopanshanensis sp. nov. from China. Mycotaxon 2019, 134, 125–137. [CrossRef]
- Chen, L.; Shi, Z.J.; Wu, C.H.; Zhao, C.L. *Gloeodontia yunnanensis* sp. nov. (Russulales, Basidiomycota) from China, evidenced by morphological characters and phylogenetic analyses. *Phytotaxa* 2020, 432, 111–118. [CrossRef]
- 93. Guan, Q.X.; Zhao, T.J.; Zhao, C.L. Morphological characters and phylogenetic analyses reveal two new species of *Peniophorella* from southern China. *Mycol. Prog.* **2020**, *19*, 397–404. [CrossRef]
- 94. Cui, B.K.; Wei, Y.L.; Dai, Y.C. Polypores from Zijin Mountain, Jiangsu Province. Mycosystema 2006, 25, 9–14.
- 95. Wei, Y.L. Species diversity, fungal flora and distribution characters of polypores in Changbai Mountains, China. *Chin. J. Appl. Ecol.* **2011**, *22*, 2711–2717.
- Gafforov, Y.; Ordynets, A.; Langer, E.; Yarasheva, M.; de Mello Gugliotta, A.; Schigel, D.; Pecoraro, L.; Zhou, Y.; Cai, L.; Zhou, L.W. Species diversity with comprehensive annotations of wood-inhabiting poroid and corticioid fungi in Uzbekistan. *Front. Microbiol.* 2020, 11, 598321. [CrossRef]
- Dai, Y.C.; Zhou, L.W.; Yang, Z.L.; Wen, H.A.; Bau, T.; Li, T.H. A revised checklist of edible fungi in China. *Mycosystema* 2010, 29, 1–21. [CrossRef]
- 98. Dai, Y.C.; Yang, Z.L. A revised checklist of medicinal fungi in China. Mycosystema 2008, 27, 801–824. [CrossRef]
- 99. Mao, X.L. The Economic Fungi in China; Science Press: Beijing, China, 1998.
- 100. Ying, J.Z.; Mao, X.L.; Ma, Q.M. Atlas of Chinese Medicinal Fungi; Science Press: Beijing, China, 1987.
- 101. Park, W.H.; Lee, H.D. Illustrated Book of Korean Medicinal Mushrooms. Kyo-Hak Publishing Co., Ltd.: Seoul, Korea, 1999.
- Saoudi, O.; Ghaouar, N. Biocatalytic characterization of free and immobilized laccase from *Trametes versicolor* in its activation zone. *Biol. Macromolocules* 2019, 128, 681–691. [CrossRef]
- 103. Riley, R.; Salamov, A.A.; Brown, D.W.; Nagy, L.G.; Floudas, D.; Held, B.W.; Levasseur, A.; Lombard, V.; Morin, E.; Otillar, R.; et al. Extensive sampling of basidiomycete genomes demonstrates inadequacy of the white-rot/brown-rot paradigm for wood decay fungi. *Proc. Natl. Acad. Sci. USA* 2014, 111, 9923–9928. [CrossRef]
- 104. Dai, Y.C. Pathogenic wood-decaying fungi on woody plants in China. Mycosystema 2012, 31, 493–509. [CrossRef]
- 105. He, M.Q.; Zhao, R.L.; Hyde, K.D.; Begerow, D.; Kemler, M.; Yurkov, A.; McKenzie, E.H.C.; Raspe, O.; Kakishima, M.; Sanchez-Ramırez, S.; et al. Notes, outline and divergence times of Basidiomycota. *Fungal Divers.* 2019, 99, 105–367. [CrossRef]
- 106. Dai, Y.C. Illustrations of Pathogenic Wood-Decaying Fungi in China; Science Press: Beijing, China, 2005.
- 107. Dai, Y.C. Illustrations of Wood-Decaying Fungi on Stored Wood or Structural Timber in China; Science Press: Beijing, China, 2009.
- 108. Dai, Y.C.; Cui, B.K. Biodiversity of Wood-Inhabiting Fungi in Hainan; Science Press: Beijing, China, 2010.
- Li, Y.; Li, T.H.; Yang, Z.L.; Bau, T.; Dai, Y.C. Atlas of Chinese Macrofungal Resources; Central China Farmer's Publishing House: Zhengzhou, China, 2015.
- 110. Liu, T.Z.; Li, G.L. Atlas of Macrofungi in Saihanwula National Nature Reserve of Inner Mongolia; Inner Mongolia Science and Technology Press: Chifeng, Mongolia, 2019.
- 111. Hawksworth, D.L.; Lücking, R. Fungal diversity revisited: 2.2 to 3.8 million species. Microbiol Spectr. 2017, 5, 4. [CrossRef]